

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 30 mg hard capsule.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains oseltamivir phosphate equivalent to 30 mg of oseltamivir.

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule

The hard capsule consists of a light yellow opaque body bearing the imprint “ROCHE” and a light yellow opaque cap bearing the imprint “30 mg”. Imprints are blue.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of influenza

In patients one year of age and older who present with symptoms typical of influenza, when influenza virus is circulating in the community. Efficacy has been demonstrated when treatment is initiated within two days of first onset of symptoms. This indication is based on clinical studies of naturally occurring influenza in which the predominant infection was influenza A (see section 5.1).

Tamiflu is indicated for the treatment of infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Prevention of influenza

- Post-exposure prevention in individuals one year of age or older following contact with a clinically diagnosed influenza case when influenza virus is circulating in the community.
- The appropriate use of Tamiflu for prevention of influenza should be determined on a case by case basis by the circumstances and the population requiring protection. In exceptional situations (e.g., in case of a mismatch between the circulating and vaccine virus strains, and a pandemic situation) seasonal prevention could be considered in individuals one year of age or older.
- Tamiflu is indicated for post-exposure prevention of influenza in infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Tamiflu is not a substitute for influenza vaccination.

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations. Decisions regarding the use of antivirals for treatment and prophylaxis should take into consideration what is known about the characteristics of the circulating influenza viruses and the impact of the disease in different geographical areas and patient populations.

Based on limited pharmacokinetic and safety data, Tamiflu can be used in infants below 12 months of age for treatment during a pandemic influenza outbreak. The treating physician should take into account the pathogenicity of the circulating strain and the underlying condition of the patient to ensure there is a potential benefit to the child.

4.2 Posology and method of administration

Tamiflu capsules and Tamiflu suspension are bioequivalent formulations. 75 mg doses can be administered as either

- one 75 mg capsule or
- one 30 mg capsule plus one 45 mg capsule or
- by administering one 30 mg dose plus one 45 mg dose of suspension.

Adults, adolescents or children (> 40 kg) who are unable to swallow capsules may receive appropriate doses of Tamiflu suspension.

Treatment of influenza

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

- For adolescents (13 to 17 years of age) and adults: The recommended oral dose is 75 mg oseltamivir twice daily for 5 days.
- For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

The following weight-adjusted dosing regimens are recommended for children 1 year of age and older:

Body Weight	Recommended dose for 5 days
≤ 15 kg	30 mg twice daily
> 15 kg to 23 kg	45 mg twice daily
> 23 kg to 40 kg	60 mg twice daily
> 40 kg	75 mg twice daily

Children weighing > 40 kg and who are able to swallow capsules may receive treatment with the adult dosage of 75 mg capsules twice daily for 5 days as an alternative to the recommended dose of Tamiflu suspension.

- For infants below 12 months of age: The recommended treatment dose for infants less than 12 months is between 2 mg/kg twice daily and 3 mg/kg twice daily during a pandemic influenza outbreak. This is based upon limited pharmacokinetic data indicating that these doses provide plasma drug exposures in the majority of patients similar to those shown to be clinically efficacious in older children and adults (see section 5.2). The following weight-adjusted dosing regimens are recommended for treatment of infants below 1 year of age:

Age	Recommended dose for 5 days
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month*	2 mg/kg twice daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention of influenza

Post-exposure prevention

- For adolescents (13 to 17 years of age) and adults: The recommended dose for prevention of influenza following close contact with an infected individual is 75 mg oseltamivir once daily for 10 days. Therapy should begin as soon as possible within two days of exposure to an infected individual.

➤ For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

The recommended post-exposure prevention dose of Tamiflu is:

Body Weight	Recommended dose for 10 days
≤ 15 kg	30 mg once daily
> 15 kg to 23 kg	45 mg once daily
> 23 kg to 40 kg	60 mg once daily
> 40 kg	75 mg once daily

Children weighing > 40 kg and who are able to swallow capsules may receive prevention with a 75 mg capsule once daily for 10 days as an alternative to the recommended dose of Tamiflu suspension.

➤ For infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children > 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month*	2 mg/kg once daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

Prevention during an influenza epidemic in the community

The recommended dose for prevention of influenza during a community outbreak is 75 mg oseltamivir once daily for up to 6 weeks.

Extemporaneous formulation

When Tamiflu powder for oral suspension is not available

During situations when commercially manufactured Tamiflu powder for oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu prepared by a pharmacist or prepared at home by parents or caregivers.

Pharmacy compounding

➤ Adults and children greater than 1 year who are unable to swallow intact capsules

This procedure describes the preparation of a 15 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (15 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed to provide a 5-day course of treatment or a 10-day course of prophylaxis for the patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (15 mg/ml) Prepared Based Upon the Patient's Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
10 to 15 kg	30 ml
> 15 to 23 kg	40 ml
> 23 to 40 kg	50 ml
> 40 kg	60 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 40 ml, 50 ml or 60 ml) of compounded suspension (15 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (15 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	29 ml
40 ml	8 capsules (600 mg)	Please use alternative capsule strength*	20 capsules (600 mg)	38.5 ml
50 ml	10 capsules (750 mg)	Please use alternative capsule strength*	25 capsules (750 mg)	48 ml
60 ml	12 capsules (900 mg)	20 capsules (900 mg)	30 capsules (900 mg)	57 ml

*No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (15 mg/ml) from Tamiflu capsules:

- Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
- Triturate the granules to a fine powder.
- Add one-third (1/3) of the specified amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) and triturate the powder until a uniform suspension is achieved.
- Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
- Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
- Repeat the rinsing (Step 5) with the remainder of the vehicle.
- Close the bottle using a child-resistant cap.
- Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
- Put an ancillary label on the bottle indicating "Shake Gently Before Use".
- Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
- Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (15 mg/ml)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature “do not store above 25 °C”.

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient’s name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension from Tamiflu Capsules for Children One Year of Age or Older

Body Weight (kg)	Dose (mg)	Volume per Dose 15 mg/ml	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
10 kg to 15 kg	30 mg	2 ml	2 ml twice daily	2 ml once daily
> 15 to 23 kg	45 mg	3 ml	3 ml twice daily	3 ml once daily
> 23 to 40 kg	60 mg	4 ml	4 ml twice daily	4 ml once daily
> 40 kg	75 mg	5 ml	5 ml twice daily	5 ml once daily

Note: This compounding procedure results in a 15 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose (2 ml, 3 ml, 4 ml or 5 ml) on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

➤ Infants less than 1 year of age

This procedure describes the preparation of a 10 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (10 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed for each patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (10 mg/ml) Prepared Based Upon the Patient’s Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
Up to 7 kg	30 ml
7 to 12 kg	45 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 45 ml) of compounded suspension (10 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (10 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	4 capsules (300 mg)	Please use alternative capsule strength*	10 capsules (300 mg)	29.5 ml
45 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	44 ml

* No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (10 mg/mL) from Tamiflu capsules:

1. Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
2. Triturate the granules to a fine powder.
3. Add one-third (1/3) of the specified amount of vehicle and triturate the powder until a uniform suspension is achieved.
4. Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
5. Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
6. Repeat the rinsing (Step 5) with the remainder of the vehicle.
7. Close the bottle using a child-resistant cap.
8. Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
9. Put an ancillary label on the bottle indicating "Shake Gently Before Use".
10. Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
11. Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (10 mg/mL)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature "do not store above 25 °C".

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient's name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants Less Than One Month of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
3 kg	0.60 ml twice daily	0.60 ml once daily
3.5 kg	0.70 ml twice daily	0.70 ml once daily
4 kg	0.80 ml twice daily	0.80 ml once daily
4.5 kg	0.90 ml twice daily	0.90 ml once daily

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants One to Twelve Months of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
4 kg	1.00 ml twice daily	1.00 ml once daily
4.5 kg	1.10 ml twice daily	1.10 ml once daily
5 kg	1.30 ml twice daily	1.30 ml once daily
5.5 kg	1.40 ml twice daily	1.40 ml once daily
6 kg	1.50 ml twice daily	1.50 ml once daily
7 kg	2.10 ml twice daily	2.10 ml once daily
8 kg	2.40 ml twice daily	2.40 ml once daily
9 kg	2.70 ml twice daily	2.70 ml once daily
≥ 10 kg	3.00 ml twice daily	3.00 ml once daily

Note: This compounding procedure results in a 10 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

Home preparation

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu (see section 3 in Package Leaflet) by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Special populations

Hepatic impairment

No dose adjustment is required either for treatment or for prevention in patients with hepatic dysfunction. No studies have been carried out in paediatric patients with hepatic disorder.

Renal impairment

Treatment of influenza: Dose adjustment is recommended for adults with severe renal impairment. Recommended doses are detailed in the table below.

Creatinine clearance	Recommended dose for treatment
> 30 (ml/min)	75 mg twice daily
> 10 to ≤ 30 (ml/min)	75 mg once daily, or 30 mg suspension twice daily, or 30 mg capsules twice daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Prevention of influenza: Dose adjustment is recommended for adults with severe renal impairment as detailed in the table below.

Creatinine clearance	Recommended dose for prevention
> 30 (ml/min)	75 mg once daily
> 10 to ≤ 30 (ml/min)	75 mg every second day, or 30 mg suspension once daily, or 30 mg capsules once daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Elderly

No dose adjustment is required, unless there is evidence of severe renal impairment.

Children

There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings and precautions for use

Oseltamivir is effective only against illness caused by influenza viruses. There is no evidence for efficacy of oseltamivir in any illness caused by agents other than influenza viruses.

No information is available regarding the safety and efficacy of oseltamivir in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalisation.

The safety and efficacy of oseltamivir in either treatment or prevention of influenza in immunocompromised patients have not been established.

Efficacy of oseltamivir in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population (see section 5.1).

Tamiflu is not a substitute for influenza vaccination. Use of Tamiflu must not affect the evaluation of individuals for annual influenza vaccination. The protection against influenza lasts only as long as Tamiflu is administered. Tamiflu should be used for the treatment and prevention of influenza only when reliable epidemiological data indicate that influenza virus is circulating in the community.

Severe renal impairment

Dose adjustment is recommended for both treatment and prevention in adults with severe renal insufficiency. There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.(see sections 4.2 and 5.2).

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic properties of oseltamivir, such as low protein binding and metabolism independent of the CYP450 and glucuronidase systems (see section 5.2), suggest that clinically significant drug interactions via these mechanisms are unlikely.

No dose adjustment is required when co-administering with probenecid in patients with normal renal function. Co-administration of probenecid, a potent inhibitor of the anionic pathway of renal tubular secretion, results in an approximate 2-fold increase in exposure to the active metabolite of oseltamivir. Oseltamivir has no kinetic interaction with amoxicillin, which is eliminated via the same pathway, suggesting that oseltamivir interaction with this pathway is weak.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely, due to the known safety margin for most of these substances, the elimination characteristics of the active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. However, care should be taken when prescribing oseltamivir in subjects when taking co-excreted agents with a narrow therapeutic margin (e.g., chlorpropamide, methotrexate, phenylbutazone).

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetyl-salicylic acid, cimetidine or with antacids (magnesium and aluminium hydroxides and calcium carbonates).

4.6 Pregnancy and lactation

While no controlled clinical trials have been conducted on the use of oseltamivir in pregnant women, there is limited data available from post-marketing and retrospective observational surveillance reports. These data in conjunction with animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal or postnatal development (see section 5.3). Pregnant women may receive Tamiflu, after considering the available safety information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the pregnant woman.

In lactating rats, oseltamivir and the active metabolite are excreted in milk. Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. Limited data demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a subtherapeutic dose to the infant. Considering this information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the lactating woman, administration of oseltamivir may be considered, where there are clear potential benefits to lactating mothers.

4.7 Effects on ability to drive and use machines

Tamiflu has no influence on the ability to drive and use machines.

4.8 Undesirable effects

The overall safety profile of Tamiflu is based on data from 2107 adult and 1032 paediatric patients treated for influenza, and on data from 2914 adult and 99 paediatric patients receiving Tamiflu for the prophylaxis of influenza in clinical trials.

In adults, the most commonly reported adverse drug reactions (ADRs) were vomiting and nausea in the treatment studies, and nausea and headache in the prevention studies. The majority of these ADRs were reported on a single occasion on either the first or second treatment day and resolved spontaneously within 1-2 days. In children, the most commonly reported adverse drug reaction was vomiting.

The ADRs listed in the tables below fall into the following categories: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). ADRs are added to the appropriate category in the tables according to the pooled analysis from clinical trials. Within each frequency grouping ADRs are presented in the order of decreasing seriousness.

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group) in Studies Investigating Tamiflu for Treatment and Prevention of Influenza in Adults and Adolescents or Through Post-Marketing Surveillance

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
Infections and infestations				
<i>Common:</i>				
Bronchitis	4 %	5 %	1 %	1 %
Bronchitis acute	1 %	1 %	0 %	< 1 %
Upper respiratory tract infections	0 %	0 %	8 %	8 %
Psychiatric disorders				
<i>Uncommon:</i>				
Hallucination ^a	< 1 %	0 %	< 1 %	0 %
Nervous system disorders				
<i>Very Common:</i>				
Headache	2 %	2 %	20 %	18 %
<i>Common:</i>				
Insomnia	1 %	1 %	1 %	1 %
<i>Uncommon:</i>				
Convulsion ^a	< 1 %	0 %	0 %	0 %
Disorders of the ear and labyrinth				
<i>Common:</i>				
Vertigo	1 %	1 %	< 1 %	< 1 %
Respiratory, thoracic and mediastinal disorders				
<i>Common:</i>				
Cough	1 %	1 %	6 %	6 %
Rhinorrhoea	< 1 %	0 %	2 %	1 %
Gastrointestinal disorders				
<i>Very Common:</i>				
Nausea ^{b,c}	11 %	7 %	8 %	4 %
<i>Common:</i>				
Vomiting ^c	8 %	3 %	2 %	1 %
Abdominal pain	2 %	2 %	2 %	2 %
Diarrhoea	6 %	8 %	3 %	3 %
Dyspepsia	1 %	1 %	2 %	2 %
Skin and subcutaneous tissue disorders				
<i>Uncommon:</i>				
Dermatitis ^a	< 1 %	< 1 %	1 %	1 %
Rash ^a	< 1 %	< 1 %	< 1 %	< 1 %
Urticaria ^a	< 1 %	< 1 %	< 1 %	< 1 %
Eczema ^a	< 1 %	0 %	< 1 %	< 1 %
General disorders				
<i>Common:</i>				
Dizziness	2 %	3 %	2 %	2 %
Fatigue	1 %	1 %	8 %	8 %
Pain	< 1 %	< 1 %	4 %	3 %

^a These are events identified during post-marketing surveillance. They were also reported in the pooled clinical studies at the incidence presented in the table above.

^b Subjects who experienced nausea alone; excludes subjects who experienced nausea in association with vomiting.

^c The difference between the placebo and oseltamivir groups was statistically significant.

Treatment and prevention of influenza in children:

The table below shows the most frequently reported ADRs from paediatric clinical trials.

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group in the treatment studies and $\geq 10\%$ in the oseltamivir group in the prophylaxis study) in Children

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Infections and infestations <i>Common:</i> Pneumonia Sinusitis Bronchitis Otitis media	2 % 2 % 2 % 9 %	3 % 3 % 2 % 11 %	0 % 0 % 2 % 1 %	0 % 0 % 0 % 2 %
Disorders of the blood and lymphatic system <i>Common:</i> Lymphadenopathy	1 %	2 %	< 1 %	0 %
Respiratory, thoracic and mediastinal disorders <i>Common:</i> Asthma (incl. aggravated) Epistaxis	4 % 3 %	4 % 3 %	0 % 1 %	1 % 1 %
Gastrointestinal disorder <i>Very Common:</i> Vomiting Diarrhoea <i>Common:</i> Nausea Abdominal pain	15 % 10 % 3 % 5 %	9 % 11 % 4 % 4 %	20 % 3 % 6 % 2 %	10 % 1 % 4 % 1 %
Disorders of the eye <i>Common:</i> Conjunctivitis	1 %	< 1 %	0 %	0 %
Disorders of the ear and labyrinth <i>Common:</i> Ear disorder ^c Tympanic membrane disorder	2 % 1 %	1 % 1 %	0 % 0 %	0 % 0 %
Skin and subcutaneous tissue disorders <i>Common:</i> Dermatitis	1 %	2 %	< 1 %	0 %

^a The prevention study did not contain a placebo arm, i.e. was an uncontrolled study.

^b Unit dose = weight-based dosing (see section 4.2).

^c Patients experienced ear ache and ear pain.

In general, the adverse event profile in children with pre-existing bronchial asthma was qualitatively similar to that of otherwise healthy children.

Further post marketing surveillance data on selected serious adverse drug reactions:

Immune system disorders

Frequency not known: hypersensitivity reactions, including anaphylactic/anaphylactoid reactions.

Psychiatric disorders and nervous system disorders

Frequency not known: influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

In patients with influenza who were receiving Tamiflu, there have been postmarketing reports of convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares), in a very few cases resulting in accidental injury or fatal outcomes. These events were reported primarily among pediatric and adolescent patients and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Eye disorders

Frequency not known: visual disturbance.

Cardiac disorders

Frequency not known: cardiac arrhythmia.

Gastrointestinal disorders

Frequency not known: gastrointestinal bleedings and hemorrhagic colitis.

Hepato-biliary disorders

Frequency not known: hepato-biliary system disorders, including hepatitis and elevated liver enzymes in patients with influenza-like illness. These cases include fatal fulminant hepatitis/hepatic failure.

Skin and subcutaneous tissue disorders

Frequency not known: severe skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme and angioneurotic oedema.

Additional information on special populations:

Infants less than one year of age

Safety information available on oseltamivir administered for treatment of influenza in infants less than one year of age from prospective and retrospective observational trials (comprising together more than 2400 infants of that age class), epidemiological databases research and postmarketing reports suggest that the safety profile in infants less than one year of age is similar to the established safety profile of children aged one year and older.

Elderly patients

There were no clinically relevant differences in the safety population of the elderly subjects who received oseltamivir or placebo compared with the adult population aged up to 65 years.

Patients with chronic cardiac and/or respiratory disease

The adverse event profile in adolescents and patients with chronic cardiac and/or respiratory disease was qualitatively similar to those of healthy young adults.

4.9 Overdose

There is no experience with overdose. However, the anticipated manifestations of acute overdose would be nausea, with or without accompanying vomiting, and dizziness. Patients should discontinue the treatment in the event of overdose. No specific antidote is known.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiviral ATC code: J05AH02

Oseltamivir phosphate is a pro-drug of the active metabolite (oseltamivir carboxylate). The active metabolite is a selective inhibitor of influenza virus neuraminidase enzymes, which are glycoproteins found on the virion surface. Viral neuraminidase enzyme activity is important both for viral entry into uninfected cells and for the release of recently formed virus particles from infected cells, and for the further spread of infectious virus in the body.

Oseltamivir carboxylate inhibits influenza A and B neuraminidases *in vitro*. Oseltamivir phosphate inhibits influenza virus infection and replication *in vitro*. Oseltamivir given orally inhibits influenza A and B virus replication and pathogenicity *in vivo* in animal models of influenza infection at antiviral exposures similar to that achieved in man with 75 mg twice daily.

Antiviral activity of oseltamivir was supported for influenza A and B by experimental challenge studies in healthy volunteers.

Neuraminidase enzyme IC₅₀ values for oseltamivir for clinically isolated influenza A ranged from 0.1 nM to 1.3 nM, and for influenza B was 2.6 nM. Higher IC₅₀ values for influenza B, up to a median of 8.5 nM, have been observed in published trials.

Reduced sensitivity of viral neuraminidase

There has been no evidence for emergence of drug resistance associated with the use of Tamiflu in clinical studies conducted to date in post-exposure (7 days), post-exposure within household groups (10 days) and seasonal (42 days) prevention of influenza.

The risk of emergence of influenza viruses with reduced susceptibility or frank resistance to oseltamivir has been examined during Roche-sponsored clinical studies. All patients who were found to carry oseltamivir-resistant virus did so transiently, cleared the virus normally and showed no clinical deterioration.

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping*	Geno- and Phenotyping*
Adults and adolescents	4/1245 (0.32%)	5/1245 (0.4%)
Children (1-12 years)	19/464 (4.1%)	25/464 (5.4%)

* Full genotyping was not performed in all studies.

The rate of emergence of resistance may be higher in the youngest age groups, and in immunosuppressed patients. Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance mutations tend to be viral sub-type specific (including those found in H5N1 variants).

Naturally occurring mutations in influenza A/H1N1 virus associated with reduced susceptibility to oseltamivir *in vitro* have been detected in patients who, based on the reported information, have not been exposed to oseltamivir. The extent of reduction in susceptibility to oseltamivir and the prevalence of such viruses appears to vary seasonally and geographically.

Treatment of influenza infection

Oseltamivir is effective only against illnesses caused by influenza virus. Statistical analyses are therefore presented only for influenza-infected subjects. In the pooled treatment study population, which included both influenza-positive and -negative subjects (ITT), primary efficacy was reduced proportional to the number of influenza-negative individuals. In the overall treatment population, influenza infection was confirmed in 67 % (range 46 % to 74 %) of the recruited patients. Of the

elderly subjects, 64 % were influenza-positive and of those with chronic cardiac and/or respiratory disease 62 % were influenza-positive. In all phase III treatment studies, patients were recruited only during the period in which influenza was circulating in the local community.

Adults and adolescents 13 years of age and older: Patients were eligible if they reported within 36 hours of onset of symptoms, had fever ≥ 37.8 °C, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache). In a pooled analysis of all influenza-positive adults and adolescents (N = 2413) enrolled into treatment studies, oseltamivir 75 mg twice daily for 5 days reduced the median duration of influenza illness by approximately one day from 5.2 days (95 % CI 4.9 – 5.5 days) in the placebo group to 4.2 days (95 % CI 4.0 – 4.4 days; $p \leq 0.0001$).

The proportion of subjects who developed specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics was reduced from 12.7 % (135/1063) in the placebo group to 8.6 % (116/1350) in the oseltamivir treated population ($p = 0.0012$).

Treatment of influenza in high risk populations: The median duration of influenza illness in elderly subjects (≥ 65 years) and in subjects with chronic cardiac and/or respiratory disease receiving oseltamivir 75 mg twice daily for 5 days was not reduced significantly. The total duration of fever was reduced by one day in the groups treated with oseltamivir. In the influenza-positive elderly, oseltamivir significantly reduced the incidence of specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics from 19 % (52/268) in the placebo group to 12 % (29/250) in the oseltamivir treated population ($p = 0.0156$).

In influenza-positive patients with chronic cardiac and/or respiratory disease, the combined incidence of lower respiratory tract complications (mainly bronchitis) treated with antibiotics was 17 % (22/133) in the placebo group and 14 % (16/118) in the oseltamivir treated population ($p = 0.5976$).

Treatment of influenza in children: In a study of otherwise healthy children (65 % influenza-positive) aged 1 to 12 years (mean age 5.3 years) who had fever (≥ 37.8 °C) plus either cough or coryza, 67 % of influenza-positive patients were infected with influenza A and 33 % with influenza B. Oseltamivir treatment, started within 48 hours of onset of symptoms, significantly reduced the time to freedom from illness (defined as the simultaneous return to normal health and activity and alleviation of fever, cough and coryza) by 1.5 days (95 % CI 0.6 – 2.2 days; $p < 0.0001$) compared to placebo. Oseltamivir reduced the incidence of acute otitis media from 26.5 % (53/200) in the placebo group to 16 % (29/183) in the oseltamivir treated children ($p = 0.013$).

A second study was completed in 334 asthmatic children aged 6 to 12 years old of which 53.6 % were influenza-positive. In the oseltamivir treated group, the median duration of illness was not reduced significantly. By day 6 (the last day of treatment) FEV₁ had increased by 10.8 % in the oseltamivir treated group compared to 4.7 % on placebo ($p = 0.0148$) in this population.

Treatment of influenza B infection: Overall, 15 % of the influenza-positive population were infected by influenza B, proportions ranging from 1 to 33 % in individual studies. The median duration of illness in influenza B infected subjects did not differ significantly between the treatment groups in individual studies. Data from 504 influenza B infected subjects were pooled across all studies for analysis. Oseltamivir reduced the time to alleviation of all symptoms by 0.7 days (95 % CI 0.1 – 1.6 days; $p = 0.022$) and the duration of fever (≥ 37.8 °C), cough and coryza by one day (95 % CI 0.4 – 1.7 days; $p < 0.001$) compared to placebo.

Prevention of influenza

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households and two seasonal prevention studies. The primary efficacy parameter for all of these studies was the incidence of laboratory-confirmed influenza. The virulence of influenza epidemics is not predictable and varies within a region and from season to season, therefore the number needed to treat (NNT) in order to prevent one case of influenza illness varies.

Post-exposure prevention: In a study in contacts (12.6 % vaccinated against influenza) of an index case of influenza, oseltamivir 75 mg once daily was started within 2 days of onset of symptoms in the index case and continued for seven days. Influenza was confirmed in 163 out of 377 index cases. Oseltamivir significantly reduced the incidence of clinical influenza illness occurring in the contacts of confirmed influenza cases from 24/200 (12 %) in the placebo group to 2/205 (1 %) in the oseltamivir group (92 % reduction [95 % CI 6 – 16; $p \leq 0.0001$]). The number needed to treat (NNT) in contacts of true influenza cases was 10 (95 % CI 9 – 12) and was 16 (95 % CI 15 – 19) in the whole population (ITT) regardless of infection status in the index case.

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households that included adults, adolescents, and children aged 1 to 12 years, both as index cases and as family contacts. The primary efficacy parameter for this study was the incidence of laboratory-confirmed clinical influenza in the households. Oseltamivir prophylaxis lasted for 10 days. In the total population, there was a reduction in the incidence of laboratory-confirmed clinical influenza in households from 20 % (27/136) in the group not receiving prevention to 7 % (10/135) in the group receiving prevention (62.7 % reduction [95 % CI 26.0 – 81.2; $p = 0.0042$]). In households of influenza-infected index cases, there was a reduction in the incidence of influenza from 26 % (23/89) in the group not receiving prevention to 11 % (9/84) in the group receiving prevention (58.5 % reduction [95 % CI 15.6 – 79.6; $p = 0.0114$]). According to subgroup analysis in children at 1 to 12 years of age, the incidence of laboratory-confirmed clinical influenza among children was significantly reduced from 19 % (21/111) in the group not receiving prevention to 7 % (7/104) in the group receiving prevention (64.4 % reduction [95 % CI 15.8 – 85.0; $p = 0.0188$]). Among children who were not already shedding virus at baseline, the incidence of laboratory-confirmed clinical influenza was reduced from 21 % (15/70) in the group not receiving prevention to 4 % (2/47) in the group receiving prevention (80.1 % reduction [95 % CI 22.0 – 94.9; $p = 0.0206$]). The NNT for the total paediatric population was 9 (95 % CI 7 – 24) and 8 (95 % CI 6, upper limit not estimable) in the whole population (ITT) and in paediatric contacts of infected index cases (ITTII), respectively.

Prevention during an influenza epidemic in the community: In a pooled analysis of two other studies conducted in unvaccinated otherwise healthy adults, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 25/519 (4.8 %) in the placebo group to 6/520 (1.2 %) in the oseltamivir group (76 % reduction [95 % CI 1.6 – 5.7; $p = 0.0006$]) during a community outbreak of influenza. The NNT in this study was 28 (95 % CI 24 – 50). A study in elderly residents of nursing homes, where 80 % of participants received vaccine in the season of the study, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 12/272 (4.4 %) in the placebo group to 1/276 (0.4 %) in the oseltamivir group (92 % reduction [95 % CI 1.5 – 6.6; $p = 0.0015$]). The NNT in this study was 25 (95 % CI 23 – 62).

Specific studies have not been conducted to assess of the reduction in the risk of complications.

5.2 Pharmacokinetic properties

Absorption

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate (pro-drug) and is extensively converted by predominantly hepatic esterases to the active metabolite (oseltamivir carboxylate). At least 75 % of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the pro-drug is less than 5 % relative to the active metabolite. Plasma concentrations of both pro-drug and active metabolite are proportional to dose and are unaffected by co-administration with food.

Distribution

The mean volume of distribution at steady state of the oseltamivir carboxylate is approximately 23 litres in humans, a volume equivalent to extracellular body fluid. Since neuraminidase activity is extracellular, oseltamivir carboxylate distributes to all sites of influenza virus spread.

The binding of the oseltamivir carboxylate to human plasma protein is negligible (approximately 3 %).

Metabolism

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. *In vitro* studies demonstrated that neither oseltamivir nor the active metabolite is a substrate for, or an inhibitor of, the major cytochrome P450 isoforms. No phase 2 conjugates of either compound have been identified *in vivo*.

Elimination

Absorbed oseltamivir is primarily (> 90 %) eliminated by conversion to oseltamivir carboxylate. It is not further metabolised and is eliminated in the urine. Peak plasma concentrations of oseltamivir carboxylate decline with a half-life of 6 to 10 hours in most subjects. The active metabolite is eliminated entirely by renal excretion. Renal clearance (18.8 l/h) exceeds glomerular filtration rate (7.5 l/h) indicating that tubular secretion occurs in addition to glomerular filtration. Less than 20 % of an oral radiolabelled dose is eliminated in faeces.

Renal impairment

Administration of 100 mg oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. For dosing, see section 4.2.

Hepatic impairment

In vitro studies have concluded that exposure to oseltamivir is not expected to be increased significantly nor is exposure to the active metabolite expected to be significantly decreased in patients with hepatic impairment (see section 4.2).

Elderly

Exposure to the active metabolite at steady state was 25 to 35 % higher in elderly (age 65 to 78 years) compared to adults less than 65 years of age given comparable doses of oseltamivir. Half-lives observed in the elderly were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage adjustments are not required for elderly patients unless there is evidence of severe renal impairment (creatinine clearance below 30 ml/min) (see section 4.2).

Children

Children 1 year of age and older: The pharmacokinetics of oseltamivir have been evaluated in single-dose pharmacokinetic studies in children aged 1 to 16 years. Multiple-dose pharmacokinetics were studied in a small number of children enrolled in a clinical efficacy study. Younger children cleared both the pro-drug and its active metabolite faster than adults, resulting in a lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in children over 12 years of age are similar to those in adults.

Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children > 1 year of age (see sections 4.1 and 4.2). There are currently no data available in infants less than 1 month of age using Tamiflu.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity and genotoxicity. Results of the conventional rodent carcinogenicity studies showed a trend towards a dose-dependent increase in the incidence of some tumours that are typical for the rodent strains used. Considering the margins of exposure in relation to

the expected exposure in the human use, these findings do not change the benefit-risk of Tamiflu in its adopted therapeutic indications.

Teratology studies have been conducted in rats and rabbits at doses of up to 1500 mg/kg/day and 500 mg/kg/day, respectively. No effects on foetal development were observed. A rat fertility study up to a dose of 1500 mg/kg/day demonstrated no adverse effects on either sex. In pre- and post-natal rat studies, prolonged parturition was noted at 1500 mg/kg/day: the safety margin between human exposure and the highest no-effect dose (500 mg/kg/day) in rats is 480-fold for oseltamivir and 44-fold for the active metabolite, respectively. Foetal exposure in the rats and rabbits was approximately 15 to 20 % of that of the mother.

In lactating rats, oseltamivir and the active metabolite are excreted in the milk. Limited data indicate that oseltamivir and the active metabolite are excreted in human milk. Extrapolation of the animal data provides estimates of 0.01 mg/day and 0.3 mg/day for the respective compounds.

A potential for skin sensitisation to oseltamivir was observed in a "maximisation" test in guinea pigs. Approximately 50 % of the animals treated with the unformulated active ingredient showed erythema after challenging the induced animals. Reversible irritancy of rabbits' eyes was detected.

Whereas very high oral single doses of oseltamivir phosphate had no effect in adult rats, such doses resulted in toxicity in juvenile 7-day-old rat pups, including death. These effects were seen at doses of 657 mg/kg and higher. At 500 mg/kg, no adverse effects were seen, including upon chronic treatment (500 mg/kg/day administered from 7 to 21 days post partum).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule core:

Pregelatinized starch (derived from maize starch)

Talc

Povidone

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell:

Gelatin

Yellow iron oxide (E172)

Red iron oxide (E172)

Titanium dioxide (E171)

Printing ink:

Shellac

Titanium dioxide (E171)

FD and C Blue 2 (indigo carmine, E132)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

7 years

6.4 Special precautions for storage

Do not store above 25 °C.

Storage of the pharmacy-compounded suspension:

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature “do not store above 25 °C”.

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

6.5 Nature and contents of container

One box contains 10 capsules in a triplex blister pack (PVC/PE/PVDC, sealed with aluminium foil).

6.6 Special precautions for disposal

No special requirements.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/02/222/003

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 20 June 2002

Date of last renewal: 20 June 2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>.

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 45 mg hard capsule.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains oseltamivir phosphate equivalent to 45 mg of oseltamivir.

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule

The hard capsule consists of a grey opaque body bearing the imprint “ROCHE” and a grey opaque cap bearing the imprint “45 mg”. Imprints are blue.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of influenza

In patients one year of age and older who present with symptoms typical of influenza, when influenza virus is circulating in the community. Efficacy has been demonstrated when treatment is initiated within two days of first onset of symptoms. This indication is based on clinical studies of naturally occurring influenza in which the predominant infection was influenza A (see section 5.1).

Tamiflu is indicated for the treatment of infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Prevention of influenza

- Post-exposure prevention in individuals one year of age or older following contact with a clinically diagnosed influenza case when influenza virus is circulating in the community.
- The appropriate use of Tamiflu for prevention of influenza should be determined on a case by case basis by the circumstances and the population requiring protection. In exceptional situations (e.g., in case of a mismatch between the circulating and vaccine virus strains, and a pandemic situation) seasonal prevention could be considered in individuals one year of age or older.
- Tamiflu is indicated for post-exposure prevention of influenza in infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Tamiflu is not a substitute for influenza vaccination.

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations. Decisions regarding the use of antivirals for treatment and prophylaxis should take into consideration what is known about the characteristics of the circulating influenza viruses and the impact of the disease in different geographical areas and patient populations.

Based on limited pharmacokinetic and safety data, Tamiflu can be used in infants below 12 months of age for treatment during a pandemic influenza outbreak. The treating physician should take into account the pathogenicity of the circulating strain and the underlying condition of the patient to ensure there is a potential benefit to the child.

4.2 Posology and method of administration

Tamiflu capsules and Tamiflu suspension are bioequivalent formulations. 75 mg doses can be administered as either

- one 75 mg capsule or
- one 30 mg capsule plus one 45 mg capsule or
- by administering one 30 mg dose plus one 45 mg dose of suspension.

Adults, adolescents or children (> 40 kg) who are unable to swallow capsules may receive appropriate doses of Tamiflu suspension.

Treatment of influenza

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

- For adolescents (13 to 17 years of age) and adults: The recommended oral dose is 75 mg oseltamivir twice daily for 5 days.
- For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

The following weight-adjusted dosing regimens are recommended for children 1 year of age and older:

Body Weight	Recommended dose for 5 days
≤ 15 kg	30 mg twice daily
> 15 kg to 23 kg	45 mg twice daily
> 23 kg to 40 kg	60 mg twice daily
> 40 kg	75 mg twice daily

Children weighing > 40 kg and who are able to swallow capsules may receive treatment with the adult dosage of 75 mg capsules twice daily for 5 days as an alternative to the recommended dose of Tamiflu suspension.

- For infants below 12 months of age: The recommended treatment dose for infants less than 12 months is between 2 mg/kg twice daily and 3 mg/kg twice daily during a pandemic influenza outbreak. This is based upon limited pharmacokinetic data indicating that these doses provide plasma drug exposures in the majority of patients similar to those shown to be clinically efficacious in older children and adults (see section 5.2). The following weight-adjusted dosing regimens are recommended for treatment of infants below 1 year of age:

Age	Recommended dose for 5 days
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month*	2 mg/kg twice daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention of influenza

Post-exposure prevention

- For adolescents (13 to 17 years of age) and adults: The recommended dose for prevention of influenza following close contact with an infected individual is 75 mg oseltamivir once daily for 10 days. Therapy should begin as soon as possible within two days of exposure to an infected individual.

➤ For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

The recommended post-exposure prevention dose of Tamiflu is:

Body Weight	Recommended dose for 10 days
≤ 15 kg	30 mg once daily
> 15 kg to 23 kg	45 mg once daily
> 23 kg to 40 kg	60 mg once daily
> 40 kg	75 mg once daily

Children weighing > 40 kg and who are able to swallow capsules may receive prevention with a 75 mg capsule once daily for 10 days as an alternative to the recommended dose of Tamiflu suspension.

➤ For infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children > 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month*	2 mg/kg once daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

Prevention during an influenza epidemic in the community

The recommended dose for prevention of influenza during a community outbreak is 75 mg oseltamivir once daily for up to 6 weeks.

Extemporaneous formulation

When Tamiflu powder for oral suspension is not available

During situations when commercially manufactured Tamiflu powder for oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu prepared by a pharmacist or prepared at home by parents or caregivers.

Pharmacy compounding

➤ Adults and children greater than 1 year who are unable to swallow intact capsules

This procedure describes the preparation of a 15 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (15 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed to provide a 5-day course of treatment or a 10-day course of prophylaxis for the patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (15 mg/ml) Prepared Based Upon the Patient's Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
10 to 15 kg	30 ml
> 15 to 23 kg	40 ml
> 23 to 40 kg	50 ml
> 40 kg	60 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 40 ml, 50 ml or 60 ml) of compounded suspension (15 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (15 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	29 ml
40 ml	8 capsules (600 mg)	Please use alternative capsule strength*	20 capsules (600 mg)	38.5 ml
50 ml	10 capsules (750 mg)	Please use alternative capsule strength*	25 capsules (750 mg)	48 ml
60 ml	12 capsules (900 mg)	20 capsules (900 mg)	30 capsules (900 mg)	57 ml

*No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (15 mg/ml) from Tamiflu capsules:

- Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
- Triturate the granules to a fine powder.
- Add one-third (1/3) of the specified amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) and triturate the powder until a uniform suspension is achieved.
- Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
- Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
- Repeat the rinsing (Step 5) with the remainder of the vehicle.
- Close the bottle using a child-resistant cap.
- Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
- Put an ancillary label on the bottle indicating "Shake Gently Before Use".
- Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
- Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (15 mg/ml)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature “do not store above 25 °C”.

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient’s name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension from Tamiflu Capsules for Children One Year of Age or Older

Body Weight (kg)	Dose (mg)	Volume per Dose 15 mg/ml	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
10 kg to 15 kg	30 mg	2 ml	2 ml twice daily	2 ml once daily
> 15 to 23 kg	45 mg	3 ml	3 ml twice daily	3 ml once daily
> 23 to 40 kg	60 mg	4 ml	4 ml twice daily	4 ml once daily
> 40 kg	75 mg	5 ml	5 ml twice daily	5 ml once daily

Note: This compounding procedure results in a 15 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose (2 ml, 3 ml, 4 ml or 5 ml) on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

➤ Infants less than 1 year of age

This procedure describes the preparation of a 10 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (10 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed for each patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (10 mg/ml) Prepared Based Upon the Patient’s Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
Up to 7 kg	30 ml
7 to 12 kg	45 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 45 ml) of compounded suspension (10 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (10 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	4 capsules (300 mg)	Please use alternative capsule strength*	10 capsules (300 mg)	29.5 ml
45 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	44 ml

* No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (10 mg/mL) from Tamiflu capsules:

- Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
- Triturate the granules to a fine powder.
- Add one-third (1/3) of the specified amount of vehicle and triturate the powder until a uniform suspension is achieved.
- Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
- Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
- Repeat the rinsing (Step 5) with the remainder of the vehicle.
- Close the bottle using a child-resistant cap.
- Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
- Put an ancillary label on the bottle indicating "Shake Gently Before Use".
- Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
- Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (10 mg/mL)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature "do not store above 25 °C".

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient's name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants Less Than One Month of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
3 kg	0.60 ml twice daily	0.60 ml once daily
3.5 kg	0.70 ml twice daily	0.70 ml once daily
4 kg	0.80 ml twice daily	0.80 ml once daily
4.5 kg	0.90 ml twice daily	0.90 ml once daily

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants One to Twelve Months of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
4 kg	1.00 ml twice daily	1.00 ml once daily
4.5 kg	1.10 ml twice daily	1.10 ml once daily
5 kg	1.30 ml twice daily	1.30 ml once daily
5.5 kg	1.40 ml twice daily	1.40 ml once daily
6 kg	1.50 ml twice daily	1.50 ml once daily
7 kg	2.10 ml twice daily	2.10 ml once daily
8 kg	2.40 ml twice daily	2.40 ml once daily
9 kg	2.70 ml twice daily	2.70 ml once daily
≥ 10 kg	3.00 ml twice daily	3.00 ml once daily

Note: This compounding procedure results in a 10 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

Home preparation

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu (see section 3 in Package Leaflet) by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Special populations

Hepatic impairment

No dose adjustment is required either for treatment or for prevention in patients with hepatic dysfunction. No studies have been carried out in paediatric patients with hepatic disorder.

Renal impairment

Treatment of influenza: Dose adjustment is recommended for adults with severe renal impairment. Recommended doses are detailed in the table below.

Creatinine clearance	Recommended dose for treatment
> 30 (ml/min)	75 mg twice daily
> 10 to ≤ 30 (ml/min)	75 mg once daily, or 30 mg suspension twice daily, or 30 mg capsules twice daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Prevention of influenza: Dose adjustment is recommended for adults with severe renal impairment as detailed in the table below.

Creatinine clearance	Recommended dose for prevention
> 30 (ml/min)	75 mg once daily
> 10 to ≤ 30 (ml/min)	75 mg every second day, or 30 mg suspension once daily, or 30 mg capsules once daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Elderly

No dose adjustment is required, unless there is evidence of severe renal impairment.

Children

There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings and precautions for use

Oseltamivir is effective only against illness caused by influenza viruses. There is no evidence for efficacy of oseltamivir in any illness caused by agents other than influenza viruses.

No information is available regarding the safety and efficacy of oseltamivir in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalisation.

The safety and efficacy of oseltamivir in either treatment or prevention of influenza in immunocompromised patients have not been established.

Efficacy of oseltamivir in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population (see section 5.1).

Tamiflu is not a substitute for influenza vaccination. Use of Tamiflu must not affect the evaluation of individuals for annual influenza vaccination. The protection against influenza lasts only as long as Tamiflu is administered. Tamiflu should be used for the treatment and prevention of influenza only when reliable epidemiological data indicate that influenza virus is circulating in the community.

Severe renal impairment

Dose adjustment is recommended for both treatment and prevention in adults with severe renal insufficiency. There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.(see sections 4.2 and 5.2).

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic properties of oseltamivir, such as low protein binding and metabolism independent of the CYP450 and glucuronidase systems (see section 5.2), suggest that clinically significant drug interactions via these mechanisms are unlikely.

No dose adjustment is required when co-administering with probenecid in patients with normal renal function. Co-administration of probenecid, a potent inhibitor of the anionic pathway of renal tubular secretion, results in an approximate 2-fold increase in exposure to the active metabolite of oseltamivir. Oseltamivir has no kinetic interaction with amoxicillin, which is eliminated via the same pathway, suggesting that oseltamivir interaction with this pathway is weak.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely, due to the known safety margin for most of these substances, the elimination characteristics of the

active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. However, care should be taken when prescribing oseltamivir in subjects when taking co-excreted agents with a narrow therapeutic margin (e.g., chlorpropamide, methotrexate, phenylbutazone).

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetyl-salicylic acid, cimetidine or with antacids (magnesium and aluminium hydroxides and calcium carbonates).

4.6 Pregnancy and lactation

While no controlled clinical trials have been conducted on the use of oseltamivir in pregnant women, there is limited data available from post-marketing and retrospective observational surveillance reports. These data in conjunction with animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal or postnatal development (see section 5.3). Pregnant women may receive Tamiflu, after considering the available safety information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the pregnant woman.

In lactating rats, oseltamivir and the active metabolite are excreted in milk. Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. Limited data demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a subtherapeutic dose to the infant. Considering this information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the lactating woman, administration of oseltamivir may be considered, where there are clear potential benefits to lactating mothers.

4.7 Effects on ability to drive and use machines

Tamiflu has no influence on the ability to drive and use machines.

4.8 Undesirable effects

The overall safety profile of Tamiflu is based on data from 2107 adult and 1032 paediatric patients treated for influenza, and on data from 2914 adult and 99 paediatric patients receiving Tamiflu for the prophylaxis of influenza in clinical trials.

In adults, the most commonly reported adverse drug reactions (ADRs) were vomiting and nausea in the treatment studies, and nausea and headache in the prevention studies. The majority of these ADRs were reported on a single occasion on either the first or second treatment day and resolved spontaneously within 1-2 days. In children, the most commonly reported adverse drug reaction was vomiting.

The ADRs listed in the tables below fall into the following categories: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). ADRs are added to the appropriate category in the tables according to the pooled analysis from clinical trials. Within each frequency grouping ADRs are presented in the order of decreasing seriousness.

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group) in Studies Investigating Tamiflu for Treatment and Prevention of Influenza in Adults and Adolescents or Through Post-Marketing Surveillance

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
Infections and infestations				
<i>Common:</i>				
Bronchitis	4 %	5 %	1 %	1 %
Bronchitis acute	1 %	1 %	0 %	< 1 %
Upper respiratory tract infections	0 %	0 %	8 %	8 %
Psychiatric disorders				
<i>Uncommon:</i>				
Hallucination ^a	< 1 %	0 %	< 1 %	0 %
Nervous system disorders				
<i>Very Common:</i>				
Headache	2 %	2 %	20 %	18 %
<i>Common:</i>				
Insomnia	1 %	1 %	1 %	1 %
<i>Uncommon:</i>				
Convulsion ^a	< 1 %	0 %	0 %	0 %
Disorders of the ear and labyrinth				
<i>Common:</i>				
Vertigo	1 %	1 %	< 1 %	< 1 %
Respiratory, thoracic and mediastinal disorders				
<i>Common:</i>				
Cough	1 %	1 %	6 %	6 %
Rhinorrhoea	< 1 %	0 %	2 %	1 %
Gastrointestinal disorders				
<i>Very Common:</i>				
Nausea ^{b,c}	11 %	7 %	8 %	4 %
<i>Common:</i>				
Vomiting ^c	8 %	3 %	2 %	1 %
Abdominal pain	2 %	2 %	2 %	2 %
Diarrhoea	6 %	8 %	3 %	3 %
Dyspepsia	1 %	1 %	2 %	2 %
Skin and subcutaneous tissue disorders				
<i>Uncommon:</i>				
Dermatitis ^a	< 1 %	< 1 %	1 %	1 %
Rash ^a	< 1 %	< 1 %	< 1 %	< 1 %
Urticaria ^a	< 1 %	< 1 %	< 1 %	< 1 %
Eczema ^a	< 1 %	0 %	< 1 %	< 1 %
General disorders				
<i>Common:</i>				
Dizziness	2 %	3 %	2 %	2 %
Fatigue	1 %	1 %	8 %	8 %
Pain	< 1 %	< 1 %	4 %	3 %

^a These are events identified during post-marketing surveillance. They were also reported in the pooled clinical studies at the incidence presented in the table above.

^b Subjects who experienced nausea alone; excludes subjects who experienced nausea in association with vomiting.

^c The difference between the placebo and oseltamivir groups was statistically significant.

Treatment and prevention of influenza in children:

The table below shows the most frequently reported ADRs from paediatric clinical trials.

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group in the treatment studies and ≥ 10 in the oseltamivir group in the prophylaxis study) in Children

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Infections and infestations <i>Common:</i> Pneumonia Sinusitis Bronchitis Otitis media	2 % 2 % 2 % 9 %	3 % 3 % 2 % 11 %	0 % 0 % 2 % 1 %	0 % 0 % 0 % 2 %
Disorders of the blood and lymphatic system <i>Common:</i> Lymphadenopathy	1 %	2 %	< 1 %	0 %
Respiratory, thoracic and mediastinal disorders <i>Common:</i> Asthma (incl. aggravated) Epistaxis	4 % 3 %	4 % 3 %	0 % 1 %	1 % 1 %
Gastrointestinal disorder <i>Very Common:</i> Vomiting Diarrhoea <i>Common:</i> Nausea Abdominal pain	15 % 10 % 3 % 5 %	9 % 11 % 4 % 4 %	20 % 3 % 6 % 2 %	10 % 1 % 4 % 1 %
Disorders of the eye <i>Common:</i> Conjunctivitis	1 %	< 1 %	0 %	0 %
Disorders of the ear and labyrinth <i>Common:</i> Ear disorder ^c Tympanic membrane disorder	2 % 1 %	1 % 1 %	0 % 0 %	0 % 0 %
Skin and subcutaneous tissue disorders <i>Common:</i> Dermatitis	1 %	2 %	< 1 %	0 %

^a The prevention study did not contain a placebo arm, i.e. was an uncontrolled study.

^b Unit dose = weight-based dosing (see section 4.2).

^c Patients experienced ear ache and ear pain.

In general, the adverse event profile in children with pre-existing bronchial asthma was qualitatively similar to that of otherwise healthy children.

Further post marketing surveillance data on selected serious adverse drug reactions:

Immune system disorders

Frequency not known: hypersensitivity reactions, including anaphylactic/anaphylactoid reactions.

Psychiatric disorders and nervous system disorders

Frequency not known: influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

In patients with influenza who were receiving Tamiflu, there have been postmarketing reports of convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares), in a very few cases resulting in accidental injury or fatal outcomes. These events were reported primarily among pediatric and adolescent patients and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Eye disorders

Frequency not known: visual disturbance.

Cardiac disorders

Frequency not known: cardiac arrhythmia.

Gastrointestinal disorders

Frequency not known: gastrointestinal bleedings and hemorrhagic colitis.

Hepato-biliary disorders

Frequency not known: hepato-biliary system disorders, including hepatitis and elevated liver enzymes in patients with influenza-like illness. These cases include fatal fulminant hepatitis/hepatic failure.

Skin and subcutaneous tissue disorders

Frequency not known: severe skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme and angioneurotic oedema.

Additional information on special populations:

Infants less than one year of age

Safety information available on oseltamivir administered for treatment of influenza in infants less than one year of age from prospective and retrospective observational trials (comprising together more than 2400 infants of that age class), epidemiological databases research and postmarketing reports suggest that the safety profile in infants less than one year of age is similar to the established safety profile of children aged one year and older.

Elderly patients

There were no clinically relevant differences in the safety population of the elderly subjects who received oseltamivir or placebo compared with the adult population aged up to 65 years.

Patients with chronic cardiac and/or respiratory disease

The adverse event profile in adolescents and patients with chronic cardiac and/or respiratory disease was qualitatively similar to those of healthy young adults.

4.9 Overdose

There is no experience with overdose. However, the anticipated manifestations of acute overdose would be nausea, with or without accompanying vomiting, and dizziness. Patients should discontinue the treatment in the event of overdose. No specific antidote is known.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiviral ATC code: J05AH02

Oseltamivir phosphate is a pro-drug of the active metabolite (oseltamivir carboxylate). The active metabolite is a selective inhibitor of influenza virus neuraminidase enzymes, which are glycoproteins found on the virion surface. Viral neuraminidase enzyme activity is important both for viral entry into uninfected cells and for the release of recently formed virus particles from infected cells, and for the further spread of infectious virus in the body.

Oseltamivir carboxylate inhibits influenza A and B neuraminidases *in vitro*. Oseltamivir phosphate inhibits influenza virus infection and replication *in vitro*. Oseltamivir given orally inhibits influenza A and B virus replication and pathogenicity *in vivo* in animal models of influenza infection at antiviral exposures similar to that achieved in man with 75 mg twice daily.

Antiviral activity of oseltamivir was supported for influenza A and B by experimental challenge studies in healthy volunteers.

Neuraminidase enzyme IC₅₀ values for oseltamivir for clinically isolated influenza A ranged from 0.1 nM to 1.3 nM, and for influenza B was 2.6 nM. Higher IC₅₀ values for influenza B, up to a median of 8.5 nM, have been observed in published trials.

Reduced sensitivity of viral neuraminidase

There has been no evidence for emergence of drug resistance associated with the use of Tamiflu in clinical studies conducted to date in post-exposure (7 days), post-exposure within household groups (10 days) and seasonal (42 days) prevention of influenza.

The risk of emergence of influenza viruses with reduced susceptibility or frank resistance to oseltamivir has been examined during Roche-sponsored clinical studies. All patients who were found to carry oseltamivir-resistant virus did so transiently, cleared the virus normally and showed no clinical deterioration.

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping*	Geno- and Phenotyping*
Adults and adolescents	4/1245 (0.32%)	5/1245 (0.4%)
Children (1-12 years)	19/464 (4.1%)	25/464 (5.4%)

* Full genotyping was not performed in all studies.

The rate of emergence of resistance may be higher in the youngest age groups, and in immunosuppressed patients. Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance mutations tend to be viral sub-type specific (including those found in H5N1 variants).

Naturally occurring mutations in influenza A/H1N1 virus associated with reduced susceptibility to oseltamivir *in vitro* have been detected in patients who, based on the reported information, have not been exposed to oseltamivir. The extent of reduction in susceptibility to oseltamivir and the prevalence of such viruses appears to vary seasonally and geographically.

Treatment of influenza infection

Oseltamivir is effective only against illnesses caused by influenza virus. Statistical analyses are therefore presented only for influenza-infected subjects. In the pooled treatment study population, which included both influenza-positive and -negative subjects (ITT), primary efficacy was reduced proportional to the number of influenza-negative individuals. In the overall treatment population, influenza infection was confirmed in 67 % (range 46 % to 74 %) of the recruited patients. Of the elderly subjects, 64 % were influenza-positive and of those with chronic cardiac and/or respiratory disease 62 % were influenza-positive. In all phase III treatment studies, patients were recruited only during the period in which influenza was circulating in the local community.

Adults and adolescents 13 years of age and older: Patients were eligible if they reported within 36 hours of onset of symptoms, had fever ≥ 37.8 °C, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache). In a pooled analysis of all influenza-positive adults and adolescents (N = 2413) enrolled into treatment studies, oseltamivir 75 mg twice daily for 5 days reduced the median duration of influenza illness by approximately one day from 5.2 days (95 % CI 4.9 – 5.5 days) in the placebo group to 4.2 days (95 % CI 4.0 – 4.4 days; $p \leq 0.0001$).

The proportion of subjects who developed specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics was reduced from 12.7 % (135/1063) in the placebo group to 8.6 % (116/1350) in the oseltamivir treated population ($p = 0.0012$).

Treatment of influenza in high risk populations: The median duration of influenza illness in elderly subjects (≥ 65 years) and in subjects with chronic cardiac and/or respiratory disease receiving oseltamivir 75 mg twice daily for 5 days was not reduced significantly. The total duration of fever was reduced by one day in the groups treated with oseltamivir. In the influenza-positive elderly, oseltamivir significantly reduced the incidence of specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics from 19 % (52/268) in the placebo group to 12 % (29/250) in the oseltamivir treated population ($p = 0.0156$).

In influenza-positive patients with chronic cardiac and/or respiratory disease, the combined incidence of lower respiratory tract complications (mainly bronchitis) treated with antibiotics was 17 % (22/133) in the placebo group and 14 % (16/118) in the oseltamivir treated population ($p = 0.5976$).

Treatment of influenza in children: In a study of otherwise healthy children (65 % influenza-positive) aged 1 to 12 years (mean age 5.3 years) who had fever (≥ 37.8 °C) plus either cough or coryza, 67 % of influenza-positive patients were infected with influenza A and 33 % with influenza B. Oseltamivir treatment, started within 48 hours of onset of symptoms, significantly reduced the time to freedom from illness (defined as the simultaneous return to normal health and activity and alleviation of fever, cough and coryza) by 1.5 days (95 % CI 0.6 – 2.2 days; $p < 0.0001$) compared to placebo. Oseltamivir reduced the incidence of acute otitis media from 26.5 % (53/200) in the placebo group to 16 % (29/183) in the oseltamivir treated children ($p = 0.013$).

A second study was completed in 334 asthmatic children aged 6 to 12 years old of which 53.6 % were influenza-positive. In the oseltamivir treated group, the median duration of illness was not reduced significantly. By day 6 (the last day of treatment) FEV₁ had increased by 10.8 % in the oseltamivir treated group compared to 4.7 % on placebo ($p = 0.0148$) in this population.

Treatment of influenza B infection: Overall, 15 % of the influenza-positive population were infected by influenza B, proportions ranging from 1 to 33 % in individual studies. The median duration of illness in influenza B infected subjects did not differ significantly between the treatment groups in individual studies. Data from 504 influenza B infected subjects were pooled across all studies for analysis. Oseltamivir reduced the time to alleviation of all symptoms by 0.7 days (95 % CI 0.1 – 1.6 days; $p = 0.022$) and the duration of fever (≥ 37.8 °C), cough and coryza by one day (95 % CI 0.4 – 1.7 days; $p < 0.001$) compared to placebo.

Prevention of influenza

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households and two seasonal prevention studies. The primary efficacy parameter for all of these studies was the incidence of laboratory-confirmed influenza. The virulence of influenza epidemics is not predictable and varies within a region and from season to season, therefore the number needed to treat (NNT) in order to prevent one case of influenza illness varies.

Post-exposure prevention: In a study in contacts (12.6 % vaccinated against influenza) of an index case of influenza, oseltamivir 75 mg once daily was started within 2 days of onset of symptoms in the index case and continued for seven days. Influenza was confirmed in 163 out of 377 index cases. Oseltamivir significantly reduced the incidence of clinical influenza illness occurring in the contacts of confirmed influenza cases from 24/200 (12 %) in the placebo group to 2/205 (1 %) in the oseltamivir group (92 % reduction [95 % CI 6 – 16; $p \leq 0.0001$]). The number needed to treat (NNT) in contacts of true influenza cases was 10 (95 % CI 9 – 12) and was 16 (95 % CI 15 – 19) in the whole population (ITT) regardless of infection status in the index case.

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households that included adults, adolescents, and children aged 1 to 12 years, both as index cases and as family contacts. The primary efficacy parameter for this study was the incidence of laboratory-confirmed clinical influenza in the households. Oseltamivir prophylaxis lasted for 10 days. In the total population, there was a reduction in the incidence of laboratory-confirmed clinical influenza in households from 20 % (27/136) in the group not receiving prevention to 7 % (10/135) in the group receiving prevention (62.7 % reduction [95 % CI 26.0 – 81.2; $p = 0.0042$]). In households of influenza-infected index cases, there was a reduction in the incidence of influenza from 26 % (23/89) in the group not receiving prevention to 11 % (9/84) in the group receiving prevention (58.5 % reduction [95 % CI 15.6 – 79.6; $p = 0.0114$]).

According to subgroup analysis in children at 1 to 12 years of age, the incidence of laboratory-confirmed clinical influenza among children was significantly reduced from 19 % (21/111) in the group not receiving prevention to 7 % (7/104) in the group receiving prevention (64.4 % reduction [95 % CI 15.8 – 85.0; $p = 0.0188$]). Among children who were not already shedding virus at baseline, the incidence of laboratory-confirmed clinical influenza was reduced from 21 % (15/70) in the group not receiving prevention to 4 % (2/47) in the group receiving prevention (80.1 % reduction [95 % CI 22.0 – 94.9; $p = 0.0206$]). The NNT for the total paediatric population was 9 (95 % CI 7 – 24) and 8 (95 % CI 6, upper limit not estimable) in the whole population (ITT) and in paediatric contacts of infected index cases (ITTII), respectively.

Prevention during an influenza epidemic in the community: In a pooled analysis of two other studies conducted in unvaccinated otherwise healthy adults, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 25/519 (4.8 %) in the placebo group to 6/520 (1.2 %) in the oseltamivir group (76 % reduction [95 % CI 1.6 – 5.7; $p = 0.0006$]) during a community outbreak of influenza. The NNT in this study was 28 (95 % CI 24 – 50).

A study in elderly residents of nursing homes, where 80 % of participants received vaccine in the season of the study, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 12/272 (4.4 %) in the placebo group to 1/276 (0.4 %) in the oseltamivir group (92 % reduction [95 % CI 1.5 – 6.6; $p = 0.0015$]). The NNT in this study was 25 (95 % CI 23 – 62).

Specific studies have not been conducted to assess of the reduction in the risk of complications.

5.2 Pharmacokinetic properties

Absorption

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate (pro-drug) and is extensively converted by predominantly hepatic esterases to the active metabolite (oseltamivir carboxylate). At least 75 % of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the pro-drug is less than 5 % relative to the active metabolite.

Plasma concentrations of both pro-drug and active metabolite are proportional to dose and are unaffected by co-administration with food.

Distribution

The mean volume of distribution at steady state of the oseltamivir carboxylate is approximately 23 litres in humans, a volume equivalent to extracellular body fluid. Since neuraminidase activity is extracellular, oseltamivir carboxylate distributes to all sites of influenza virus spread.

The binding of the oseltamivir carboxylate to human plasma protein is negligible (approximately 3 %).

Metabolism

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. *In vitro* studies demonstrated that neither oseltamivir nor the active metabolite is a substrate for, or an inhibitor of, the major cytochrome P450 isoforms. No phase 2 conjugates of either compound have been identified *in vivo*.

Elimination

Absorbed oseltamivir is primarily (> 90 %) eliminated by conversion to oseltamivir carboxylate. It is not further metabolised and is eliminated in the urine. Peak plasma concentrations of oseltamivir carboxylate decline with a half-life of 6 to 10 hours in most subjects. The active metabolite is eliminated entirely by renal excretion. Renal clearance (18.8 l/h) exceeds glomerular filtration rate (7.5 l/h) indicating that tubular secretion occurs in addition to glomerular filtration. Less than 20 % of an oral radiolabelled dose is eliminated in faeces.

Renal impairment

Administration of 100 mg oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. For dosing, see section 4.2.

Hepatic impairment

In vitro studies have concluded that exposure to oseltamivir is not expected to be increased significantly nor is exposure to the active metabolite expected to be significantly decreased in patients with hepatic impairment (see section 4.2).

Elderly

Exposure to the active metabolite at steady state was 25 to 35 % higher in elderly (age 65 to 78 years) compared to adults less than 65 years of age given comparable doses of oseltamivir. Half-lives observed in the elderly were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage adjustments are not required for elderly patients unless there is evidence of severe renal impairment (creatinine clearance below 30 ml/min) (see section 4.2).

Children

Children one year of age and older: The pharmacokinetics of oseltamivir have been evaluated in single-dose pharmacokinetic studies in children aged 1 to 16 years. Multiple-dose pharmacokinetics were studied in a small number of children enrolled in a clinical efficacy study. Younger children cleared both the pro-drug and its active metabolite faster than adults, resulting in a lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in children over 12 years of age are similar to those in adults.

Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children > 1 year of age (see sections 4.1 and 4.2). There are currently no data available in patients less than 1 month of age using Tamiflu.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity and genotoxicity. Results of the conventional rodent carcinogenicity studies showed a trend towards a dose-dependent increase in the incidence of some tumours that are typical for the rodent strains used. Considering the margins of exposure in relation to the expected exposure in the human use, these findings do not change the benefit-risk of Tamiflu in its adopted therapeutic indications.

Teratology studies have been conducted in rats and rabbits at doses of up to 1500 mg/kg/day and 500 mg/kg/day, respectively. No effects on foetal development were observed. A rat fertility study up to a dose of 1500 mg/kg/day demonstrated no adverse effects on either sex. In pre- and post-natal rat studies, prolonged parturition was noted at 1500 mg/kg/day: the safety margin between human exposure and the highest no-effect dose (500 mg/kg/day) in rats is 480-fold for oseltamivir and 44-fold for the active metabolite, respectively. Foetal exposure in the rats and rabbits was approximately 15 to 20 % of that of the mother.

In lactating rats, oseltamivir and the active metabolite are excreted in the milk. Limited data indicate that oseltamivir and the active metabolite are excreted in human milk. Extrapolation of the animal data provides estimates of 0.01 mg/day and 0.3 mg/day for the respective compounds.

A potential for skin sensitisation to oseltamivir was observed in a "maximisation" test in guinea pigs. Approximately 50 % of the animals treated with the unformulated active ingredient showed erythema after challenging the induced animals. Reversible irritancy of rabbits' eyes was detected.

Whereas very high oral single doses of oseltamivir phosphate had no effect in adult rats, such doses resulted in toxicity in juvenile 7-day-old rat pups, including death. These effects were seen at doses of 657 mg/kg and higher. At 500 mg/kg, no adverse effects were seen, including upon chronic treatment (500 mg/kg/day administered from 7 to 21 days post partum).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule core:

Pregelatinized starch (derived from maize starch)

Talc

Povidone

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell:

Gelatin

Black iron oxide (E172)

Titanium dioxide (E171)

Printing ink:

Shellac

Titanium dioxide (E171)

FD and C Blue 2 (indigo carmine, E132)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

7 years

6.4 Special precautions for storage

Do not store above 25 °C.

Storage of the pharmacy-compounded suspension:

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature “do not store above 25 °C”.

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

6.5 Nature and contents of container

One box contains 10 capsules in a triplex blister pack (PVC/PE/PVDC, sealed with aluminium foil).

6.6 Special precautions for disposal

No special requirements.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/02/222/004

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 20 June 2002

Date of last renewal: 20 June 2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>.

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 75 mg hard capsule.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains oseltamivir phosphate equivalent to 75 mg of oseltamivir.
For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule

The hard capsule consists of a grey opaque body bearing the imprint “ROCHE” and a light yellow opaque cap bearing the imprint “75 mg”. Imprints are blue.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of influenza

In patients one year of age and older who present with symptoms typical of influenza, when influenza virus is circulating in the community. Efficacy has been demonstrated when treatment is initiated within two days of first onset of symptoms. This indication is based on clinical studies of naturally occurring influenza in which the predominant infection was influenza A (see section 5.1).

Tamiflu is indicated for the treatment of infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Prevention of influenza

- Post-exposure prevention in individuals one year of age or older following contact with a clinically diagnosed influenza case when influenza virus is circulating in the community.
- The appropriate use of Tamiflu for prevention of influenza should be determined on a case by case basis by the circumstances and the population requiring protection. In exceptional situations (e.g., in case of a mismatch between the circulating and vaccine virus strains, and a pandemic situation) seasonal prevention could be considered in individuals one year of age or older.
- Tamiflu is indicated for post-exposure prevention of influenza in infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Tamiflu is not a substitute for influenza vaccination.

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations. Decisions regarding the use of antivirals for treatment and prophylaxis should take into consideration what is known about the characteristics of the circulating influenza viruses and the impact of the disease in different geographical areas and patient populations.

Based on limited pharmacokinetic and safety data, Tamiflu can be used in infants below 12 months of age for treatment during a pandemic influenza outbreak. The treating physician should take into account the pathogenicity of the circulating strain and the underlying condition of the patient to ensure there is a potential benefit to the child.

4.2 Posology and method of administration

Tamiflu capsules and Tamiflu suspension are bioequivalent formulations. 75 mg doses can be administered as either

- one 75 mg capsule or
- one 30 mg capsule plus one 45 mg capsule or
- by administering one 30 mg dose plus one 45 mg dose of suspension.

Adults, adolescents or children (> 40 kg) who are unable to swallow capsules may receive appropriate doses of Tamiflu suspension.

Treatment of influenza

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

- For adolescents (13 to 17 years of age) and adults: The recommended oral dose is 75 mg oseltamivir twice daily for 5 days.
- For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

For recommended treatment dose of Tamiflu for infants older than 1 year of age and for children 2 to 12 years of age, see SmPC of Tamiflu suspension and Tamiflu 30 and 45 mg capsules.

Children weighing > 40 kg and who are able to swallow capsules may receive treatment with the adult dosage of 75 mg capsules twice daily for 5 days as an alternative to the recommended dose of Tamiflu suspension or Tamiflu 30 mg and 45 mg capsules.

- For infants below 12 months of age: The recommended treatment dose for infants less than 12 months is between 2 mg/kg twice daily and 3 mg/kg twice daily during a pandemic influenza outbreak. This is based upon limited pharmacokinetic data indicating that these doses provide plasma drug exposures in the majority of patients similar to those shown to be clinically efficacious in older children and adults (see section 5.2). The following weight-adjusted dosing regimens are recommended for treatment of infants below 1 year of age:

Age	Recommended dose for 5 days
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month*	2 mg/kg twice daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention of influenza

Post-exposure prevention

- For adolescents (13 to 17 years of age) and adults: The recommended dose for prevention of influenza following close contact with an infected individual is 75 mg oseltamivir once daily for 10 days. Therapy should begin as soon as possible within two days of exposure to an infected individual.
- For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules and oral suspension are available.

For recommended post-exposure prevention dose of Tamiflu for infants older than 1 year of age and for children of 2 to 12 years of age, see SmPC of Tamiflu suspension and Tamiflu 30 mg and 45 mg capsules.

Children weighing > 40 kg and who are able to swallow capsules may receive prevention with a 75 mg capsule once daily for 10 days as an alternative to the recommended dose of Tamiflu suspension or Tamiflu 30 mg and 45 mg capsules.

➤ For infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children > 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month*	2 mg/kg once daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

Prevention during an influenza epidemic in the community

The recommended dose for prevention of influenza during a community outbreak is 75 mg oseltamivir once daily for up to 6 weeks.

Extemporaneous formulation

When Tamiflu powder for oral suspension is not available

During situations when commercially manufactured Tamiflu powder for oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu prepared by a pharmacist or prepared at home by parents or caregivers.

Pharmacy compounding

➤ Adults and children greater than 1 year who are unable to swallow intact capsules

This procedure describes the preparation of a 15 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (15 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed to provide a 5-day course of treatment or a 10-day course of prophylaxis for the patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (15 mg/ml) Prepared Based Upon the Patient's Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
10 to 15 kg	30 ml
> 15 to 23 kg	40 ml
> 23 to 40 kg	50 ml
> 40 kg	60 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 40 ml, 50 ml or 60 ml) of compounded suspension (15 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (15 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	29 ml
40 ml	8 capsules (600 mg)	Please use alternative capsule strength*	20 capsules (600 mg)	38.5 ml
50 ml	10 capsules (750 mg)	Please use alternative capsule strength*	25 capsules (750 mg)	48 ml
60 ml	12 capsules (900 mg)	20 capsules (900 mg)	30 capsules (900 mg)	57 ml

* No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (15 mg/ml) from Tamiflu capsules:

- Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
- Triturate the granules to a fine powder.
- Add one-third (1/3) of the specified amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) and triturate the powder until a uniform suspension is achieved.
- Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
- Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
- Repeat the rinsing (Step 5) with the remainder of the vehicle.
- Close the bottle using a child-resistant cap.
- Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
- Put an ancillary label on the bottle indicating "Shake Gently Before Use".
- Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
- Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (15 mg/ml)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature "do not store above 25 °C".

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient's name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension from Tamiflu Capsules for Children One Year of Age or Older

Body Weight (kg)	Dose (mg)	Volume per Dose 15 mg/ml	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
10 kg to 15 kg	30 mg	2 ml	2 ml twice daily	2 ml once daily
> 15 to 23 kg	45 mg	3 ml	3 ml twice daily	3 ml once daily
> 23 to 40 kg	60 mg	4 ml	4 ml twice daily	4 ml once daily
> 40 kg	75 mg	5 ml	5 ml twice daily	5 ml once daily

Note: This compounding procedure results in a 15 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose (2 ml, 3 ml, 4 ml or 5 ml) on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

➤ *Infants less than 1 year of age*

This procedure describes the preparation of a 10 mg/ml solution that will provide one patient with enough medication for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a suspension (10 mg/ml) from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.1% w/v sodium benzoate added as a preservative.

First, calculate the Total Volume needed to be compounded and dispensed for each patient. The Total Volume required is determined by the weight of the patient according to the recommendation in the table below:

Volume of Compounded Suspension (10 mg/ml) Prepared Based Upon the Patient's Weight

Body Weight (kg)	Total Volume to Compound per Patient Weight (ml)
Up to 7 kg	30 ml
7 to 12 kg	45 ml

Second, determine the number of capsules and the amount of vehicle (water containing 0.1% w/v sodium benzoate added as a preservative) that is needed to prepare the Total Volume (calculated from the table above: 30 ml, 45 ml) of compounded suspension (10 mg/ml) as shown in the table below:

Number of Capsules and Amount of Vehicle Needed to Prepare the Total Volume of a Compounded Suspension (10 mg/ml)

Total Volume of Compounded Suspension to be Prepared	Required Number of Tamiflu Capsules (mg of oseltamivir)			Required Volume of Vehicle
	75 mg	45 mg	30 mg	
30 ml	4 capsules (300 mg)	Please use alternative capsule strength*	10 capsules (300 mg)	29.5 ml
45 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	44 ml

* No integral number of capsules can be used to achieve the target concentration; therefore, please use either the 30 mg or 75 mg capsules.

Third, follow the procedure below for compounding the suspension (10 mg/mL) from Tamiflu capsules:

1. Carefully separate the capsule body and cap and transfer the contents of the required number of Tamiflu capsules into a clean mortar.
2. Triturate the granules to a fine powder.
3. Add one-third (1/3) of the specified amount of vehicle and triturate the powder until a uniform suspension is achieved.
4. Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
5. Add another one-third (1/3) of the vehicle to the mortar, rinse the pestle and mortar by a triturating motion and transfer the vehicle into the bottle.
6. Repeat the rinsing (Step 5) with the remainder of the vehicle.
7. Close the bottle using a child-resistant cap.
8. Shake well to completely dissolve the active drug and to ensure homogeneous distribution of the dissolved drug in the resulting suspension.
(Note: Undissolved residue may be visible but is comprised of inert ingredients of Tamiflu capsules, which are insoluble. However, the active drug, oseltamivir phosphate, readily dissolves in the specified vehicle and therefore forms a uniform solution.)
9. Put an ancillary label on the bottle indicating "Shake Gently Before Use".
10. Instruct the parent or caregiver that after the patient has completed the full course of therapy any remaining solution must be discarded. It is recommended that this information be provided by affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
11. Place an appropriate expiration date label according to storage condition (see below).

Storage of the pharmacy-compounded suspension (10 mg/mL)

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature "do not store above 25 °C".

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

Place a pharmacy label on the bottle that includes the patient's name, dosing instructions, use by date, drug name and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants Less Than One Month of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
3 kg	0.60 ml twice daily	0.60 ml once daily
3.5 kg	0.70 ml twice daily	0.70 ml once daily
4 kg	0.80 ml twice daily	0.80 ml once daily
4.5 kg	0.90 ml twice daily	0.90 ml once daily

Dosing Chart for Pharmacy-Compounded Suspension (10 mg/ml) from Tamiflu Capsules for Infants One to Twelve Months of Age

Body Weight (rounded to the nearest 0.5 kg)	Treatment Dose (for 5 days)	Prophylaxis Dose (for 10 days)
4 kg	1.00 ml twice daily	1.00 ml once daily
4.5 kg	1.10 ml twice daily	1.10 ml once daily
5 kg	1.30 ml twice daily	1.30 ml once daily
5.5 kg	1.40 ml twice daily	1.40 ml once daily
6 kg	1.50 ml twice daily	1.50 ml once daily
7 kg	2.10 ml twice daily	2.10 ml once daily
8 kg	2.40 ml twice daily	2.40 ml once daily
9 kg	2.70 ml twice daily	2.70 ml once daily
≥ 10 kg	3.00 ml twice daily	3.00 ml once daily

Note: This compounding procedure results in a 10 mg/ml suspension, which is different from the commercially available Tamiflu powder for oral suspension.

Dispense the suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

Home preparation

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu (see section 3 in Package Leaflet) by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Special populations

Hepatic impairment

No dose adjustment is required either for treatment or for prevention in patients with hepatic dysfunction. No studies have been carried out in paediatric patients with hepatic disorder.

Renal impairment

Treatment of influenza: Dose adjustment is recommended for adults with severe renal impairment. Recommended doses are detailed in the table below.

Creatinine clearance	Recommended dose for treatment
> 30 (ml/min)	75 mg twice daily
> 10 to ≤ 30 (ml/min)	75 mg once daily, or 30 mg suspension twice daily, or 30 mg capsules twice daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Prevention of influenza: Dose adjustment is recommended for adults with severe renal impairment as detailed in the table below.

Creatinine clearance	Recommended dose for prevention
> 30 (ml/min)	75 mg once daily
> 10 to ≤ 30 (ml/min)	75 mg every second day, or 30 mg suspension once daily, or 30 mg capsules once daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Elderly

No dose adjustment is required, unless there is evidence of severe renal impairment.

Children

There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings and precautions for use

Oseltamivir is effective only against illness caused by influenza viruses. There is no evidence for efficacy of oseltamivir in any illness caused by agents other than influenza viruses.

No information is available regarding the safety and efficacy of oseltamivir in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalisation.

The safety and efficacy of oseltamivir in either treatment or prevention of influenza in immunocompromised patients have not been established.

Efficacy of oseltamivir in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population (see section 5.1).

Tamiflu is not a substitute for influenza vaccination. Use of Tamiflu must not affect the evaluation of individuals for annual influenza vaccination. The protection against influenza lasts only as long as Tamiflu is administered. Tamiflu should be used for the treatment and prevention of influenza only when reliable epidemiological data indicate that influenza virus is circulating in the community.

Severe renal impairment

Dose adjustment is recommended for both treatment and prevention in adults with severe renal insufficiency. There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.(see sections 4.2 and 5.2).

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic properties of oseltamivir, such as low protein binding and metabolism independent of the CYP450 and glucuronidase systems (see section 5.2), suggest that clinically significant drug interactions via these mechanisms are unlikely.

No dose adjustment is required when co-administering with probenecid in patients with normal renal function. Co-administration of probenecid, a potent inhibitor of the anionic pathway of renal tubular secretion, results in an approximate 2-fold increase in exposure to the active metabolite of oseltamivir. Oseltamivir has no kinetic interaction with amoxicillin, which is eliminated via the same pathway, suggesting that oseltamivir interaction with this pathway is weak.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely, due to the known safety margin for most of these substances, the elimination characteristics of the active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. However, care should be taken when prescribing oseltamivir in subjects when taking co-excreted agents with a narrow therapeutic margin (e.g., chlorpropamide, methotrexate, phenylbutazone).

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetyl-salicylic acid, cimetidine or with antacids (magnesium and aluminium hydroxides and calcium carbonates).

4.6 Pregnancy and lactation

While no controlled clinical trials have been conducted on the use of oseltamivir in pregnant women, there is limited data available from post-marketing and retrospective observational surveillance reports. These data in conjunction with animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal or postnatal development (see section 5.3). Pregnant women may receive Tamiflu, after considering the available safety information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the pregnant woman.

In lactating rats, oseltamivir and the active metabolite are excreted in milk. Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. Limited data demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a subtherapeutic dose to the infant. Considering this information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the lactating woman, administration of oseltamivir may be considered, where there are clear potential benefits to lactating mothers.

4.7 Effects on ability to drive and use machines

Tamiflu has no influence on the ability to drive and use machines.

4.8 Undesirable effects

The overall safety profile of Tamiflu is based on data from 2107 adult and 1032 paediatric patients treated for influenza, and on data from 2914 adult and 99 paediatric patients receiving Tamiflu for the prophylaxis of influenza in clinical trials.

In adults, the most commonly reported adverse drug reactions (ADRs) were vomiting and nausea in the treatment studies, and nausea and headache in the prevention studies. The majority of these ADRs were reported on a single occasion on either the first or second treatment day and resolved spontaneously within 1-2 days. In children, the most commonly reported adverse drug reaction was vomiting.

The ADRs listed in the tables below fall into the following categories: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). ADRs are added to the appropriate category in the tables according to the pooled analysis from clinical trials. Within each frequency grouping ADRs are presented in the order of decreasing seriousness.

Treatment and prevention of influenza in adults and adolescents:

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group) in Studies Investigating Tamiflu for Treatment and Prevention of Influenza in Adults and Adolescents or Through Post-Marketing Surveillance

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
Infections and infestations				
Common:				
Bronchitis	4 %	5 %	1 %	1 %
Bronchitis acute	1 %	1 %	0 %	< 1 %
Upper respiratory tract infections	0 %	0 %	8 %	8 %
Psychiatric disorders				
Uncommon:				
Hallucination ^a	< 1 %	0 %	< 1 %	0 %
Nervous system disorders				
Very Common:				
Headache	2 %	2 %	20 %	18 %
Common:				
Insomnia	1 %	1 %	1 %	1 %
Uncommon:				
Convulsion ^a	< 1 %	0 %	0 %	0 %
Disorders of the ear and labyrinth				
Common:				
Vertigo	1 %	1 %	< 1 %	< 1 %

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
Respiratory, thoracic and mediastinal disorders <i>Common:</i> Cough Rhinorrhoea	1 % < 1 %	1 % 0 %	6 % 2 %	6 % 1 %
Gastrointestinal disorders <i>Very Common:</i> Nausea ^{b,c} <i>Common:</i> Vomiting ^c Abdominal pain Diarrhoea Dyspepsia	11 % 8 % 2 % 6 % 1 %	7 % 3 % 2 % 8 % 1 %	8 % 2 % 2 % 3 % 2 %	4 % 1 % 2 % 3 % 2 %
Skin and subcutaneous tissue disorders <i>Uncommon:</i> Dermatitis ^a Rash ^a Urticaria ^a Eczema ^a	< 1 % < 1 % < 1 % < 1 %	< 1 % < 1 % < 1 % 0 %	1 % < 1 % < 1 % < 1 %	1 % < 1 % < 1 % < 1 %
General disorders <i>Common:</i> Dizziness Fatigue Pain	2 % 1 % < 1 %	3 % 1 % < 1 %	2 % 8 % 4 %	2 % 8 % 3 %

^a These are events identified during post-marketing surveillance. They were also reported in the pooled clinical studies at the incidence presented in the table above.

^b Subjects who experienced nausea alone; excludes subjects who experienced nausea in association with vomiting.

^c The difference between the placebo and oseltamivir groups was statistically significant.

Treatment and prevention of influenza in children:

The table below shows the most frequently reported ADRs from paediatric clinical trials.

Most Frequent Adverse Drug Reactions (≥ 1 % in the oseltamivir group in the treatment studies and ≥ 10 % in the oseltamivir group in the prophylaxis study) in Children

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Infections and infestations <i>Common:</i> Pneumonia Sinusitis Bronchitis Otitis media	2 % 2 % 2 % 9 %	3 % 3 % 2 % 11 %	0 % 0 % 2 % 1 %	0 % 0 % 0 % 2 %

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Disorders of the blood and lymphatic system <i>Common:</i> Lymphadenopathy	1 %	2 %	< 1 %	0 %
Respiratory, thoracic and mediastinal disorders <i>Common:</i> Asthma (incl. aggravated) Epistaxis	4 % 3 %	4 % 3 %	0 % 1 %	1 % 1 %
Gastrointestinal disorder <i>Very Common:</i> Vomiting Diarrhoea <i>Common:</i> Nausea Abdominal pain	15 % 10 % 3 % 5 %	9 % 11 % 4 % 4 %	20 % 3 % 6 % 2 %	10 % 1 % 4 % 1 %
Disorders of the eye <i>Common:</i> Conjunctivitis	1 %	< 1 %	0 %	0 %
Disorders of the ear and labyrinth <i>Common:</i> Ear disorder ^c Tympanic membrane disorder	2 % 1 %	1 % 1 %	0 % 0 %	0 % 0 %
Skin and subcutaneous tissue disorders <i>Common:</i> Dermatitis	1 %	2 %	< 1 %	0 %

^a The prevention study did not contain a placebo arm, i.e. was an uncontrolled study.

^b Unit dose = weight-based dosing (see section 4.2).

^c Patients experienced ear ache and ear pain.

In general, the adverse event profile in children with pre-existing bronchial asthma was qualitatively similar to that of otherwise healthy children.

Further post marketing surveillance data on selected serious adverse drug reactions:

Immune system disorders

Frequency not known: hypersensitivity reactions, including anaphylactic/anaphylactoid reactions.

Psychiatric disorders and nervous system disorders

Frequency not known: influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

In patients with influenza who were receiving Tamiflu, there have been postmarketing reports of convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares), in a very few cases resulting in accidental injury or fatal outcomes. These events were reported primarily among pediatric and adolescent patients and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Eye disorders

Frequency not known: visual disturbance.

Cardiac disorders

Frequency not known: cardiac arrhythmia.

Gastrointestinal disorders

Frequency not known: gastrointestinal bleedings and hemorrhagic colitis.

Hepato-biliary disorders

Frequency not known: hepato-biliary system disorders, including hepatitis and elevated liver enzymes in patients with influenza-like illness. These cases include fatal fulminant hepatitis/hepatic failure.

Skin and subcutaneous tissue disorders

Frequency not known: severe skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme and angioneurotic oedema.

Additional information on special populations:

Infants less than one year of age

Safety information available on oseltamivir administered for treatment of influenza in infants less than one year of age from prospective and retrospective observational trials (comprising together more than 2400 infants of that age class), epidemiological databases research and postmarketing reports suggest that the safety profile in infants less than one year of age is similar to the established safety profile of children aged one year and older.

Elderly patients

There were no clinically relevant differences in the safety population of the elderly subjects who received oseltamivir or placebo compared with the adult population aged up to 65 years.

Patients with chronic cardiac and/or respiratory disease

The adverse event profile in adolescents and patients with chronic cardiac and/or respiratory disease was qualitatively similar to those of healthy young adults.

4.9 Overdose

There is no experience with overdose. However, the anticipated manifestations of acute overdose would be nausea, with or without accompanying vomiting, and dizziness. Patients should discontinue the treatment in the event of overdose. No specific antidote is known.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiviral ATC code: J05AH02

Oseltamivir phosphate is a pro-drug of the active metabolite (oseltamivir carboxylate). The active metabolite is a selective inhibitor of influenza virus neuraminidase enzymes, which are glycoproteins found on the virion surface. Viral neuraminidase enzyme activity is important both for viral entry into uninfected cells and for the release of recently formed virus particles from infected cells, and for the further spread of infectious virus in the body.

Oseltamivir carboxylate inhibits influenza A and B neuraminidases *in vitro*. Oseltamivir phosphate inhibits influenza virus infection and replication *in vitro*. Oseltamivir given orally inhibits influenza A

and B virus replication and pathogenicity *in vivo* in animal models of influenza infection at antiviral exposures similar to that achieved in man with 75 mg twice daily.

Antiviral activity of oseltamivir was supported for influenza A and B by experimental challenge studies in healthy volunteers.

Neuraminidase enzyme IC₅₀ values for oseltamivir for clinically isolated influenza A ranged from 0.1 nM to 1.3 nM, and for influenza B was 2.6 nM. Higher IC₅₀ values for influenza B, up to a median of 8.5 nM, have been observed in published trials.

Reduced sensitivity of viral neuraminidase

There has been no evidence for emergence of drug resistance associated with the use of Tamiflu in clinical studies conducted to date in post-exposure (7 days), post-exposure within household groups (10 days) and seasonal (42 days) prevention of influenza.

The risk of emergence of influenza viruses with reduced susceptibility or frank resistance to oseltamivir has been examined during Roche-sponsored clinical studies. All patients who were found to carry oseltamivir-resistant virus did so transiently, cleared the virus normally and showed no clinical deterioration.

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping*	Geno- and Phenotyping*
Adults and adolescents	4/1245 (0.32%)	5/1245 (0.4%)
Children (1-12 years)	19/464 (4.1%)	25/464 (5.4%)

* Full genotyping was not performed in all studies.

The rate of emergence of resistance may be higher in the youngest age groups, and in immunosuppressed patients. Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance mutations tend to be viral sub-type specific (including those found in H5N1 variants).

Naturally occurring mutations in influenza A/H1N1 virus associated with reduced susceptibility to oseltamivir *in vitro* have been detected in patients who, based on the reported information, have not been exposed to oseltamivir. The extent of reduction in susceptibility to oseltamivir and the prevalence of such viruses appears to vary seasonally and geographically.

Treatment of influenza infection

Oseltamivir is effective only against illnesses caused by influenza virus. Statistical analyses are therefore presented only for influenza-infected subjects. In the pooled treatment study population, which included both influenza-positive and -negative subjects (ITT), primary efficacy was reduced proportional to the number of influenza-negative individuals. In the overall treatment population, influenza infection was confirmed in 67 % (range 46 % to 74 %) of the recruited patients. Of the elderly subjects, 64 % were influenza-positive and of those with chronic cardiac and/or respiratory disease 62 % were influenza-positive. In all phase III treatment studies, patients were recruited only during the period in which influenza was circulating in the local community.

Adults and adolescents 13 years of age and older: Patients were eligible if they reported within 36 hours of onset of symptoms, had fever ≥ 37.8 °C, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache). In a pooled analysis of all influenza-positive adults and adolescents (N = 2413) enrolled into treatment studies, oseltamivir 75 mg twice daily for 5 days reduced the median duration of influenza illness by approximately one day from 5.2 days (95 % CI 4.9 – 5.5 days) in the placebo group to 4.2 days (95 % CI 4.0 – 4.4 days; $p \leq 0.0001$).

The proportion of subjects who developed specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics was reduced from 12.7 % (135/1063) in the placebo group to 8.6 % (116/1350) in the oseltamivir treated population ($p = 0.0012$).

Treatment of influenza in high risk populations: The median duration of influenza illness in elderly subjects (≥ 65 years) and in subjects with chronic cardiac and/or respiratory disease receiving oseltamivir 75 mg twice daily for 5 days was not reduced significantly. The total duration of fever was reduced by one day in the groups treated with oseltamivir. In the influenza-positive elderly, oseltamivir significantly reduced the incidence of specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics from 19 % (52/268) in the placebo group to 12 % (29/250) in the oseltamivir treated population ($p = 0.0156$).

In influenza-positive patients with chronic cardiac and/or respiratory disease, the combined incidence of lower respiratory tract complications (mainly bronchitis) treated with antibiotics was 17 % (22/133) in the placebo group and 14 % (16/118) in the oseltamivir treated population ($p = 0.5976$).

Treatment of influenza in children: In a study of otherwise healthy children (65 % influenza-positive) aged 1 to 12 years (mean age 5.3 years) who had fever (≥ 37.8 °C) plus either cough or coryza, 67 % of influenza-positive patients were infected with influenza A and 33 % with influenza B. Oseltamivir treatment, started within 48 hours of onset of symptoms, significantly reduced the time to freedom from illness (defined as the simultaneous return to normal health and activity and alleviation of fever, cough and coryza) by 1.5 days (95 % CI 0.6 – 2.2 days; $p < 0.0001$) compared to placebo. Oseltamivir reduced the incidence of acute otitis media from 26.5 % (53/200) in the placebo group to 16 % (29/183) in the oseltamivir treated children ($p = 0.013$).

A second study was completed in 334 asthmatic children aged 6 to 12 years old of which 53.6 % were influenza-positive. In the oseltamivir treated group, the median duration of illness was not reduced significantly. By day 6 (the last day of treatment) FEV₁ had increased by 10.8 % in the oseltamivir treated group compared to 4.7 % on placebo ($p = 0.0148$) in this population.

Treatment of influenza B infection: Overall, 15 % of the influenza-positive population were infected by influenza B, proportions ranging from 1 to 33 % in individual studies. The median duration of illness in influenza B infected subjects did not differ significantly between the treatment groups in individual studies. Data from 504 influenza B infected subjects were pooled across all studies for analysis. Oseltamivir reduced the time to alleviation of all symptoms by 0.7 days (95 % CI 0.1 – 1.6 days; $p = 0.022$) and the duration of fever (≥ 37.8 °C), cough and coryza by one day (95 % CI 0.4 – 1.7 days; $p < 0.001$) compared to placebo.

Prevention of influenza

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households and two seasonal prevention studies. The primary efficacy parameter for all of these studies was the incidence of laboratory-confirmed influenza. The virulence of influenza epidemics is not predictable and varies within a region and from season to season, therefore the number needed to treat (NNT) in order to prevent one case of influenza illness varies.

Post-exposure prevention: In a study in contacts (12.6 % vaccinated against influenza) of an index case of influenza, oseltamivir 75 mg once daily was started within 2 days of onset of symptoms in the index case and continued for seven days. Influenza was confirmed in 163 out of 377 index cases. Oseltamivir significantly reduced the incidence of clinical influenza illness occurring in the contacts of confirmed influenza cases from 24/200 (12 %) in the placebo group to 2/205 (1 %) in the oseltamivir group (92 % reduction [95 % CI 6 – 16; $p \leq 0.0001$]). The number needed to treat (NNT) in contacts of true influenza cases was 10 (95 % CI 9 – 12) and was 16 (95 % CI 15 – 19) in the whole population (ITT) regardless of infection status in the index case.

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households that included adults, adolescents, and children aged

1 to 12 years, both as index cases and as family contacts. The primary efficacy parameter for this study was the incidence of laboratory-confirmed clinical influenza in the households. Oseltamivir prophylaxis lasted for 10 days. In the total population, there was a reduction in the incidence of laboratory-confirmed clinical influenza in households from 20 % (27/136) in the group not receiving prevention to 7 % (10/135) in the group receiving prevention (62.7 % reduction [95 % CI 26.0 – 81.2; $p = 0.0042$]). In households of influenza-infected index cases, there was a reduction in the incidence of influenza from 26 % (23/89) in the group not receiving prevention to 11 % (9/84) in the group receiving prevention (58.5 % reduction [95 % CI 15.6 – 79.6; $p = 0.0114$]).

According to subgroup analysis in children at 1 to 12 years of age, the incidence of laboratory-confirmed clinical influenza among children was significantly reduced from 19 % (21/111) in the group not receiving prevention to 7 % (7/104) in the group receiving prevention (64.4 % reduction [95 % CI 15.8 – 85.0; $p = 0.0188$]). Among children who were not already shedding virus at baseline, the incidence of laboratory-confirmed clinical influenza was reduced from 21 % (15/70) in the group not receiving prevention to 4 % (2/47) in the group receiving prevention (80.1 % reduction [95 % CI 22.0 – 94.9; $p = 0.0206$]). The NNT for the total paediatric population was 9 (95 % CI 7 – 24) and 8 (95 % CI 6, upper limit not estimable) in the whole population (ITT) and in paediatric contacts of infected index cases (ITTII), respectively.

Prevention during an influenza epidemic in the community: In a pooled analysis of two other studies conducted in unvaccinated otherwise healthy adults, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 25/519 (4.8 %) in the placebo group to 6/520 (1.2 %) in the oseltamivir group (76 % reduction [95 % CI 1.6 – 5.7; $p = 0.0006$]) during a community outbreak of influenza. The NNT in this study was 28 (95 % CI 24 – 50).

A study in elderly residents of nursing homes, where 80 % of participants received vaccine in the season of the study, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 12/272 (4.4 %) in the placebo group to 1/276 (0.4 %) in the oseltamivir group (92 % reduction [95 % CI 1.5 – 6.6; $p = 0.0015$]). The NNT in this study was 25 (95 % CI 23 – 62).

Specific studies have not been conducted to assess of the reduction in the risk of complications.

5.2 Pharmacokinetic properties

Absorption

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate (pro-drug) and is extensively converted by predominantly hepatic esterases to the active metabolite (oseltamivir carboxylate). At least 75 % of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the pro-drug is less than 5 % relative to the active metabolite. Plasma concentrations of both pro-drug and active metabolite are proportional to dose and are unaffected by co-administration with food.

Distribution

The mean volume of distribution at steady state of the oseltamivir carboxylate is approximately 23 litres in humans, a volume equivalent to extracellular body fluid. Since neuraminidase activity is extracellular, oseltamivir carboxylate distributes to all sites of influenza virus spread.

The binding of the oseltamivir carboxylate to human plasma protein is negligible (approximately 3 %).

Metabolism

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. *In vitro* studies demonstrated that neither oseltamivir nor the active metabolite is a substrate for, or an inhibitor of, the major cytochrome P450 isoforms. No phase 2 conjugates of either compound have been identified *in vivo*.

Elimination

Absorbed oseltamivir is primarily (> 90 %) eliminated by conversion to oseltamivir carboxylate. It is not further metabolised and is eliminated in the urine. Peak plasma concentrations of oseltamivir

carboxylate decline with a half-life of 6 to 10 hours in most subjects. The active metabolite is eliminated entirely by renal excretion. Renal clearance (18.8 l/h) exceeds glomerular filtration rate (7.5 l/h) indicating that tubular secretion occurs in addition to glomerular filtration. Less than 20 % of an oral radiolabelled dose is eliminated in faeces.

Renal impairment

Administration of 100 mg oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. For dosing, see section 4.2.

Hepatic impairment

In vitro studies have concluded that exposure to oseltamivir is not expected to be increased significantly nor is exposure to the active metabolite expected to be significantly decreased in patients with hepatic impairment (see section 4.2).

Elderly

Exposure to the active metabolite at steady state was 25 to 35 % higher in elderly (age 65 to 78 years) compared to adults less than 65 years of age given comparable doses of oseltamivir. Half-lives observed in the elderly were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage adjustments are not required for elderly patients unless there is evidence of severe renal impairment (creatinine clearance below 30 ml/min) (see section 4.2).

Children

Children 1 year of age and older: The pharmacokinetics of oseltamivir have been evaluated in single-dose pharmacokinetic studies in children aged 1 to 16 years. Multiple-dose pharmacokinetics were studied in a small number of children enrolled in a clinical efficacy study. Younger children cleared both the pro-drug and its active metabolite faster than adults, resulting in a lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in children over 12 years of age are similar to those in adults.

Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children > 1 year of age (see sections 4.1 and 4.2). There are currently no data available in infants less than 1 month of age using Tamiflu.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity and genotoxicity. Results of the conventional rodent carcinogenicity studies showed a trend towards a dose-dependent increase in the incidence of some tumours that are typical for the rodent strains used. Considering the margins of exposure in relation to the expected exposure in the human use, these findings do not change the benefit-risk of Tamiflu in its adopted therapeutic indications.

Teratology studies have been conducted in rats and rabbits at doses of up to 1500 mg/kg/day and 500 mg/kg/day, respectively. No effects on foetal development were observed. A rat fertility study up to a dose of 1500 mg/kg/day demonstrated no adverse effects on either sex. In pre- and post-natal rat studies, prolonged parturition was noted at 1500 mg/kg/day: the safety margin between human exposure and the highest no-effect dose (500 mg/kg/day) in rats is 480-fold for oseltamivir and 44-fold for the active metabolite, respectively. Foetal exposure in the rats and rabbits was approximately 15 to 20 % of that of the mother.

In lactating rats, oseltamivir and the active metabolite are excreted in the milk. Limited data indicate that oseltamivir and the active metabolite are excreted in human milk. Extrapolation of the animal data provides estimates of 0.01 mg/day and 0.3 mg/day for the respective compounds.

A potential for skin sensitisation to oseltamivir was observed in a "maximisation" test in guinea pigs. Approximately 50 % of the animals treated with the unformulated active ingredient showed erythema after challenging the induced animals. Reversible irritancy of rabbits' eyes was detected.

Whereas very high oral single doses of oseltamivir phosphate had no effect in adult rats, such doses resulted in toxicity in juvenile 7-day-old rat pups, including death. These effects were seen at doses of 657 mg/kg and higher. At 500 mg/kg, no adverse effects were seen, including upon chronic treatment (500 mg/kg/day administered from 7 to 21 days post partum).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule core:

Pregelatinized starch (derived from maize starch)

Talc

Povidone

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell:

Gelatin

Yellow iron oxide (E172)

Red iron oxide (E172)

Black iron oxide (E172)

Titanium dioxide (E171)

Printing ink:

Shellac

Titanium dioxide (E171)

FD and C Blue 2 (indigo carmine, E132)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

7 years

6.4 Special precautions for storage

Do not store above 25 °C.

Storage of the pharmacy-compounded suspension:

Room temperature storage conditions: Stable for 3 weeks (21 days) when stored at room temperature "do not store above 25 °C".

Refrigerated storage conditions: Stable for 6 weeks when stored at 2 °C - 8 °C.

6.5 Nature and contents of container

One box contains 10 capsules in a triplex blister pack (PVC/PE/PVDC, sealed with aluminium foil).

6.6 Special precautions for disposal

No special requirements.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/02/222/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 20 June 2002

Date of last renewal: 20 June 2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>.

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 12 mg/ml powder for oral suspension.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 g of powder for oral suspension contains oseltamivir phosphate equivalent to 30 mg of oseltamivir. After reconstitution, each ml of suspension contains 12 mg oseltamivir. One bottle of reconstituted suspension (75 ml) contains 900 mg of active substance (oseltamivir).

A bottle of 30 g Tamiflu powder for oral suspension contains 25.713 g of sorbitol. One dose of 45 mg oseltamivir administered twice daily delivers 2.6 g of sorbitol.

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for oral suspension

The powder is a granulate or clumped granulate with a white to light yellow colour.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of influenza

In patients one year of age and older who present with symptoms typical of influenza, when influenza virus is circulating in the community. Efficacy has been demonstrated when treatment is initiated within two days of first onset of symptoms. This indication is based on clinical studies of naturally occurring influenza in which the predominant infection was influenza A (see section 5.1).

Tamiflu is indicated for the treatment of infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Prevention of influenza

- Post-exposure prevention in individuals one year of age or older following contact with a clinically diagnosed influenza case when influenza virus is circulating in the community.
- The appropriate use of Tamiflu for prevention of influenza should be determined on a case by case basis by the circumstances and the population requiring protection. In exceptional situations (e.g., in case of a mismatch between the circulating and vaccine virus strains, and a pandemic situation) seasonal prevention could be considered in individuals one year of age or older.
- Tamiflu is indicated for post-exposure prevention of influenza in infants below 12 months of age during a pandemic influenza outbreak (see section 5.2).

Tamiflu is not a substitute for influenza vaccination.

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations. Decisions regarding the use of antivirals for treatment and prophylaxis should take into consideration what is known about the characteristics of the circulating influenza viruses and the impact of the disease in different geographical areas and patient populations.

Based on limited pharmacokinetic and safety data, Tamiflu can be used in infants below 12 months of age for treatment during a pandemic influenza outbreak. The treating physician should take into account the pathogenicity of the circulating strain and the underlying condition of the patient to ensure there is a potential benefit to the child.

4.2 Posology and method of administration

Tamiflu suspension and Tamiflu capsules are bioequivalent formulations. 75 mg doses can be administered as either

- one 75 mg capsule or
- one 30 mg capsule plus one 45 mg capsule or
- by administering one 30 mg dose plus one 45 mg dose of suspension.

Adults, adolescents or children (> 40 kg) who are able to swallow capsules may receive appropriate doses of Tamiflu capsules.

Treatment of influenza

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

➤ For adolescents (13 to 17 years of age) and adults: The recommended oral dose is 75 mg oseltamivir twice daily for 5 days.

➤ For infants older than 1 year of age and for children 2 to 12 years of age: The recommended dose of Tamiflu oral suspension is indicated in the table below. Tamiflu 30 mg and 45 mg capsules are available as an alternative to the recommended dose of Tamiflu suspension.

The following weight-adjusted dosing regimens are recommended for children 1 year of age and older:

Body Weight	Recommended dose for 5 days
≤ 15 kg	30 mg twice daily
> 15 kg to 23 kg	45 mg twice daily
> 23 kg to 40 kg	60 mg twice daily
> 40 kg	75 mg twice daily

For dosing, an oral dispenser with 30 mg, 45 mg and 60 mg graduations is provided in the box. For accurate dosing, the oral dispenser supplied should be used exclusively.

Children weighing > 40 kg and who are able to swallow capsules may receive treatment with the adult dosage of 75 mg capsules twice daily for 5 days as an alternative to the recommended dose of Tamiflu suspension.

➤ For infants below 12 months of age: The recommended treatment dose for infants less than 12 months is between 2 mg/kg twice daily and 3 mg/kg twice daily during a pandemic influenza outbreak. This is based upon limited pharmacokinetic data indicating that these doses provide plasma drug exposures in the majority of patients similar to those shown to be clinically efficacious in older children and adults (see section 5.2). The following weight-adjusted dosing regimens are recommended for treatment of infants below 1 year of age:

Age	Recommended dose for 5 days
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month*	2 mg/kg twice daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention of influenza

Post-exposure prevention

➤ For adolescents (13 to 17 years of age) and adults: The recommended dose for prevention of influenza following close contact with an infected individual is 75 mg oseltamivir once daily for 10 days. Therapy should begin as soon as possible within two days of exposure to an infected individual.

➤ For infants older than 1 year of age and for children 2 to 12 years of age: Tamiflu 30 mg and 45 mg capsules are available as an alternative to the recommended dose of Tamiflu suspension.

The recommended post-exposure prevention dose of Tamiflu is:

Body Weight	Recommended dose for 10 days
≤ 15 kg	30 mg once daily
> 15 kg to 23 kg	45 mg once daily
> 23 kg to 40 kg	60 mg once daily
> 40 kg	75 mg once daily

For dosing, an oral dispenser with 30 mg, 45 mg and 60 mg graduations is provided in the box. For accurate dosing, the oral dispenser supplied should be used exclusively.

It is recommended that Tamiflu powder for oral suspension be constituted by a pharmacist prior to dispensing to the patient (see section 6.6).

Children weighing > 40 kg and who are able to swallow capsules may receive prophylaxis with a 75 mg capsule once daily for 10 days as an alternative to the recommended dose of Tamiflu suspension.

➤ For infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children > 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month*	2 mg/kg once daily

* There is no data available regarding the administration of Tamiflu to infants less than one month of age.

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

Prevention during an influenza epidemic in the community

The recommended dose for prevention of influenza during a community outbreak is 75 mg oseltamivir once daily for up to 6 weeks.

Special populations

Hepatic impairment

No dose adjustment is required either for treatment or for prevention in patients with hepatic dysfunction. No studies have been carried out in paediatric patients with hepatic disorder.

Renal impairment

Treatment of influenza: Dose adjustment is recommended for adults with severe renal impairment. Recommended doses are detailed in the table below.

Creatinine clearance	Recommended dose for treatment
> 30 (ml/min)	75 mg twice daily
> 10 to ≤ 30 (ml/min)	75 mg once daily, or 30 mg suspension twice daily, or 30 mg capsules twice daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Prevention of influenza: Dose adjustment is recommended for adults with severe renal impairment as detailed in the table below.

Creatinine clearance	Recommended dose for prevention
> 30 (ml/min)	75 mg once daily
> 10 to ≤ 30 (ml/min)	75 mg every second day, or 30 mg suspension once daily, or 30 mg capsules once daily
≤ 10 (ml/min)	Not recommended
dialysis patients	Not recommended

Elderly

No dose adjustment is required, unless there is evidence of severe renal impairment.

Children

There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings and precautions for use

Oseltamivir is effective only against illness caused by influenza viruses. There is no evidence for efficacy of oseltamivir in any illness caused by agents other than influenza viruses.

No information is available regarding the safety and efficacy of oseltamivir in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalisation.

The safety and efficacy of oseltamivir in either treatment or prevention of influenza in immunocompromised patients have not been established.

Efficacy of oseltamivir in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population (see section 5.1).

Tamiflu is not a substitute for influenza vaccination. Use of Tamiflu must not affect the evaluation of individuals for annual influenza vaccination. The protection against influenza lasts only as long as Tamiflu is administered. Tamiflu should be used for the treatment and prevention of influenza only when reliable epidemiological data indicate that influenza virus is circulating in the community.

Severe renal impairment

Dose adjustment is recommended for both treatment and prevention in adults with severe renal insufficiency. There is insufficient clinical data available in children with renal impairment to be able to make any dosing recommendation (see sections 4.2 and 5.2).

This medicinal product contains sorbitol. Patients with rare hereditary problems of fructose intolerance should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic properties of oseltamivir, such as low protein binding and metabolism independent of the CYP450 and glucuronidase systems (see section 5.2), suggest that clinically significant drug interactions via these mechanisms are unlikely.

No dose adjustment is required when co-administering with probenecid in patients with normal renal function. Co-administration of probenecid, a potent inhibitor of the anionic pathway of renal tubular secretion, results in an approximate 2-fold increase in exposure to the active metabolite of oseltamivir. Oseltamivir has no kinetic interaction with amoxicillin, which is eliminated via the same pathway, suggesting that oseltamivir interaction with this pathway is weak.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely, due to the known safety margin for most of these substances, the elimination characteristics of the active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. However, care should be taken when prescribing oseltamivir in subjects when taking co-excreted agents with a narrow therapeutic margin (e.g., chlorpropamide, methotrexate, phenylbutazone).

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetyl-salicylic acid, cimetidine or with antacids (magnesium and aluminium hydroxides and calcium carbonates).

4.6 Pregnancy and lactation

While no controlled clinical trials have been conducted on the use of oseltamivir in pregnant women, there is limited data available from post-marketing and retrospective observational surveillance reports. These data in conjunction with animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal or postnatal development (see section 5.3). Pregnant women may receive Tamiflu, after considering the available safety information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the pregnant woman.

In lactating rats, oseltamivir and the active metabolite are excreted in milk. Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. Limited data demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a subtherapeutic dose to the infant. Considering this information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the lactating woman, administration of oseltamivir may be considered, where there are clear potential benefits to lactating mothers.

4.7 Effects on ability to drive and use machines

Tamiflu has no influence on the ability to drive and use machines.

4.8 Undesirable effects

The overall safety profile of Tamiflu is based on data from 2107 adult and 1032 paediatric patients treated for influenza, and on data from 2914 adult and 99 paediatric patients receiving Tamiflu for the prophylaxis of influenza in clinical trials.

In adults, the most commonly reported adverse drug reactions (ADRs) were vomiting and nausea in the treatment studies, and nausea and headache in the prevention studies. The majority of these ADRs were reported on a single occasion on either the first or second treatment day and resolved

spontaneously within 1-2 days. In children, the most commonly reported adverse drug reaction was vomiting.

The ADRs listed in the tables below fall into the following categories: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). ADRs are added to the appropriate category in the tables according to the pooled analysis from clinical trials. Within each frequency grouping ADRs are presented in the order of decreasing seriousness.

Treatment and prevention of influenza in adults and adolescents:

Most Frequent Adverse Drug Reactions ($\geq 1\%$ in the oseltamivir group) in Studies Investigating Tamiflu for Treatment and Prevention of Influenza in Adults and Adolescents or Through Post-Marketing Surveillance

System Organ Class (SOC) Frequency Category Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
Infections and infestations				
<i>Common:</i>				
Bronchitis	4 %	5 %	1 %	1 %
Bronchitis acute	1 %	1 %	0 %	< 1 %
Upper respiratory tract infections	0 %	0 %	8 %	8 %
Psychiatric disorders				
<i>Uncommon:</i>				
Hallucination ^a	< 1 %	0 %	< 1 %	0 %
Nervous system disorders				
<i>Very Common:</i>				
Headache	2 %	2 %	20 %	18 %
<i>Common:</i>				
Insomnia	1 %	1 %	1 %	1 %
<i>Uncommon:</i>				
Convulsion ^a	< 1 %	0 %	0 %	0 %
Disorders of the ear and labyrinth				
<i>Common:</i>				
Vertigo	1 %	1 %	< 1 %	< 1 %
Respiratory, thoracic and mediastinal disorders				
<i>Common:</i>				
Cough	1 %	1 %	6 %	6 %
Rhinorrhoea	< 1 %	0 %	2 %	1 %
Gastrointestinal disorders				
<i>Very Common:</i>				
Nausea ^{b,c}	11 %	7 %	8 %	4 %
<i>Common:</i>				
Vomiting ^c	8 %	3 %	2 %	1 %
Abdominal pain	2 %	2 %	2 %	2 %
Diarrhoea	6 %	8 %	3 %	3 %
Dyspepsia	1 %	1 %	2 %	2 %
Skin and subcutaneous tissue disorders				
<i>Uncommon:</i>				
Dermatitis ^a	< 1 %	< 1 %	1 %	1 %
Rash ^a	< 1 %	< 1 %	< 1 %	< 1 %
Urticaria ^a	< 1 %	< 1 %	< 1 %	< 1 %
Eczema ^a	< 1 %	0 %	< 1 %	< 1 %

System Organ Class (SOC) <i>Frequency Category</i> Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Prevention	
	Oseltamivir 75 mg bid (n = 1057)	Placebo (n = 1050)	Oseltamivir 75 mg od (n = 1480)	Placebo (n = 1434)
General disorders				
<i>Common:</i>				
Dizziness	2 %	3 %	2 %	2 %
Fatigue	1 %	1 %	8 %	8 %
Pain	< 1 %	< 1 %	4 %	3 %

^a These are events identified during post-marketing surveillance. They were also reported in the pooled clinical studies the incidence presented in the table above.

^b .Subjects who experienced nausea alone; excludes subjects who experienced nausea in association with vomiting.

^c The difference between the placebo and oseltamivir groups was statistically significant.

Treatment and prevention of influenza in children:

The table below shows the most frequently reported ADRs from paediatric clinical trials.

Most Frequent Adverse Drug Reactions (≥ 1 % in the oseltamivir group in the treatment studies and ≥ 10 % in the oseltamivir group in the prophylaxis study) in Children

System Organ Class (SOC) <i>Frequency Category</i> Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Infections and infestations				
<i>Common:</i>				
Pneumonia	2 %	3 %	0 %	0 %
Sinusitis	2 %	3 %	0 %	0 %
Bronchitis	2 %	2 %	2 %	0 %
Otitis media	9 %	11 %	1 %	2 %
Disorders of the blood and lymphatic system				
<i>Common:</i>				
Lymphadenopathy	1 %	2 %	< 1 %	0 %
Respiratory, thoracic and mediastinal disorders				
<i>Common:</i>				
Asthma (incl. aggravated)	4 %	4 %	0 %	1 %
Epistaxis	3 %	3 %	1 %	1 %
Gastrointestinal disorder				
<i>Very Common:</i>				
Vomiting	15 %	9 %	20 %	10 %
Diarrhoea	10 %	11 %	3 %	1 %
<i>Common:</i>				
Nausea	3 %	4 %	6 %	4 %
Abdominal pain	5 %	4 %	2 %	1 %
Disorders of the eye				
<i>Common:</i>				
Conjunctivitis	1 %	< 1 %	0 %	0 %
Disorders of the ear and labyrinth				
<i>Common:</i>				
Ear disorder ^c	2 %	1 %	0 %	0 %
Tympanic membrane disorder	1 %	1 %	0 %	0 %

System Organ Class (SOC) <i>Frequency Category</i> Adverse Drug Reaction	Percentage of Patients Experiencing the ADR			
	Treatment		Treatment	Prevention ^a
	Oseltamivir 2 mg/kg bid (n = 515)	Placebo (n = 517)	Oseltamivir 30 to 75 mg ^b (n = 158)	Oseltamivir 30 to 75 mg ^b (n = 99)
Skin and subcutaneous tissue disorders <i>Common:</i> Dermatitis	1 %	2 %	< 1 %	0 %

^a The prevention study did not contain a placebo arm, i.e. was an uncontrolled study.

^b Unit dose = weight-based dosing (see section 4.2).

^c Patients experienced ear ache and ear pain.

In general, the adverse event profile in children with pre-existing bronchial asthma was qualitatively similar to that of otherwise healthy children.

Further post marketing surveillance data on selected serious adverse drug reactions:

Immune system disorders

Frequency not known: hypersensitivity reactions, including anaphylactic/anaphylactoid reactions.

Psychiatric disorders and nervous system disorders

Frequency not known: influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

In patients with influenza who were receiving Tamiflu, there have been postmarketing reports of convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares), in a very few cases resulting in accidental injury or fatal outcomes. These events were reported primarily among pediatric and adolescent patients and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Eye disorders

Frequency not known: visual disturbance.

Cardiac disorders

Frequency not known: cardiac arrhythmia.

Gastrointestinal disorders

Frequency not known: gastrointestinal bleedings and hemorrhagic colitis.

Hepato-biliary disorders

Frequency not known: hepato-biliary system disorders, including hepatitis and elevated liver enzymes in patients with influenza-like illness. These cases include fatal fulminant hepatitis/hepatic failure.

Skin and subcutaneous tissue disorders

Frequency not known: severe skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme and angioneurotic oedema.

Additional information on special populations:

Infants less than one year of age

Safety information available on oseltamivir administered for treatment of influenza in infants less than one year of age from prospective and retrospective observational trials (comprising together more than

2400 infants of that age class), epidemiological databases research and postmarketing reports suggest that the safety profile in infants less than one year of age is similar to the established safety profile of children aged one year and older.

Elderly patients

There were no clinically relevant differences in the safety population of the elderly subjects who received oseltamivir or placebo compared with the adult population aged up to 65 years.

Patients with chronic cardiac and/or respiratory disease

The adverse event profile in adolescents and patients with chronic cardiac and/or respiratory disease was qualitatively similar to those of healthy young adults.

4.9 Overdose

There is no experience with overdose. However, the anticipated manifestations of acute overdose would be nausea, with or without accompanying vomiting, and dizziness. Patients should discontinue the treatment in the event of overdose. No specific antidote is known.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiviral ATC code: J05AH02

Oseltamivir phosphate is a pro-drug of the active metabolite (oseltamivir carboxylate). The active metabolite is a selective inhibitor of influenza virus neuraminidase enzymes, which are glycoproteins found on the virion surface. Viral neuraminidase enzyme activity is important both for viral entry into uninfected cells and for the release of recently formed virus particles from infected cells, and for the further spread of infectious virus in the body.

Oseltamivir carboxylate inhibits influenza A and B neuraminidases *in vitro*. Oseltamivir phosphate inhibits influenza virus infection and replication *in vitro*. Oseltamivir given orally inhibits influenza A and B virus replication and pathogenicity *in vivo* in animal models of influenza infection at antiviral exposures similar to that achieved in man with 75 mg twice daily.

Antiviral activity of oseltamivir was supported for influenza A and B by experimental challenge studies in healthy volunteers.

Neuraminidase enzyme IC₅₀ values for oseltamivir for clinically isolated influenza A ranged from 0.1 nM to 1.3 nM, and for influenza B was 2.6 nM. Higher IC₅₀ values for influenza B, up to a median of 8.5 nM, have been observed in published trials.

Reduced sensitivity of viral neuraminidase

There has been no evidence for emergence of drug resistance associated with the use of Tamiflu in clinical studies conducted to date in post-exposure (7 days), post-exposure within household groups (10 days) and seasonal (42 days) prevention of influenza.

The risk of emergence of influenza viruses with reduced susceptibility or frank resistance to oseltamivir has been examined during Roche-sponsored clinical studies. All patients who were found to carry oseltamivir-resistant virus did so transiently, cleared the virus normally and showed no clinical deterioration.

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping*	Geno- and Phenotyping*
Adults and adolescents	4/1245 (0.32%)	5/1245 (0.4%)
Children (1-12 years)	19/464 (4.1%)	25/464 (5.4%)

* Full genotyping was not performed in all studies.

The rate of emergence of resistance may be higher in the youngest age groups, and in immunosuppressed patients. Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance mutations tend to be viral sub-type specific (including those found in H5N1 variants).

Naturally occurring mutations in influenza A/H1N1 virus associated with reduced susceptibility to oseltamivir *in vitro* have been detected in patients who, based on the reported information, have not been exposed to oseltamivir. The extent of reduction in susceptibility to oseltamivir and the prevalence of such viruses appears to vary seasonally and geographically.

Treatment of influenza infection

Oseltamivir is effective only against illnesses caused by influenza virus. Statistical analyses are therefore presented only for influenza-infected subjects. In the pooled treatment study population, which included both influenza-positive and -negative subjects (ITT), primary efficacy was reduced proportional to the number of influenza-negative individuals. In the overall treatment population, influenza infection was confirmed in 67 % (range 46 % to 74 %) of the recruited patients. Of the elderly subjects, 64 % were influenza-positive and of those with chronic cardiac and/or respiratory disease 62 % were influenza-positive. In all phase III treatment studies, patients were recruited only during the period in which influenza was circulating in the local community.

Adults and adolescents 13 years of age and older: Patients were eligible if they reported within 36 hours of onset of symptoms, had fever ≥ 37.8 °C, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache). In a pooled analysis of all influenza-positive adults and adolescents (N = 2413) enrolled into treatment studies, oseltamivir 75 mg twice daily for 5 days reduced the median duration of influenza illness by approximately one day from 5.2 days (95 % CI 4.9 – 5.5 days) in the placebo group to 4.2 days (95 % CI 4.0 – 4.4 days; $p \leq 0.0001$).

The proportion of subjects who developed specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics was reduced from 12.7 % (135/1063) in the placebo group to 8.6 % (116/1350) in the oseltamivir treated population ($p = 0.0012$).

Treatment of influenza in high risk populations: The median duration of influenza illness in elderly subjects (≥ 65 years) and in subjects with chronic cardiac and/or respiratory disease receiving oseltamivir 75 mg twice daily for 5 days was not reduced significantly. The total duration of fever was reduced by one day in the groups treated with oseltamivir. In the influenza-positive elderly, oseltamivir significantly reduced the incidence of specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics from 19 % (52/268) in the placebo group to 12 % (29/250) in the oseltamivir treated population ($p = 0.0156$).

In influenza-positive patients with chronic cardiac and/or respiratory disease, the combined incidence of lower respiratory tract complications (mainly bronchitis) treated with antibiotics was 17 % (22/133) in the placebo group and 14 % (16/118) in the oseltamivir treated population ($p = 0.5976$).

Treatment of influenza in children: In a study of otherwise healthy children (65 % influenza-positive) aged 1 to 12 years (mean age 5.3 years) who had fever (≥ 37.8 °C) plus either cough or coryza, 67 % of influenza-positive patients were infected with influenza A and 33 % with influenza B. Oseltamivir treatment, started within 48 hours of onset of symptoms, significantly reduced the duration of time to freedom from illness (defined as the simultaneous return to normal health and activity and alleviation of fever, cough and coryza) by 1.5 days (95 % CI 0.6 – 2.2 days; $p < 0.0001$) compared to placebo.

Oseltamivir reduced the incidence of acute otitis media from 26.5 % (53/200) in the placebo group to 16 % (29/183) in the oseltamivir treated children ($p = 0.013$).

A second study was completed in 334 asthmatic children aged 6 to 12 years old of which 53.6 % were influenza-positive. In the oseltamivir treated group, the median duration of illness was not reduced significantly. By day 6 (the last day of treatment) FEV₁ had increased by 10.8 % in the oseltamivir treated group compared to 4.7 % on placebo ($p = 0.0148$) in this population.

Treatment of influenza B infection: Overall, 15 % of the influenza-positive population were infected by influenza B, proportions ranging from 1 to 33 % in individual studies. The median duration of illness in influenza B infected subjects did not differ significantly between the treatment groups in individual studies. Data from 504 influenza B infected subjects were pooled across all studies for analysis. Oseltamivir reduced the time to alleviation of all symptoms by 0.7 days (95 % CI 0.1 – 1.6 days; $p = 0.022$) and the duration of fever (≥ 37.8 °C), cough and coryza by one day (95 % CI 0.4 – 1.7 days; $p < 0.001$) compared to placebo.

Prevention of influenza

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households and two seasonal prevention studies. The primary efficacy parameter for all of these studies was the incidence of laboratory-confirmed influenza. The virulence of influenza epidemics is not predictable and varies within a region and from season to season, therefore the number needed to treat (NNT) in order to prevent one case of influenza illness varies.

Post-exposure prevention: In a study in contacts (12.6 % vaccinated against influenza) of an index case of influenza, oseltamivir 75 mg once daily was started within 2 days of onset of symptoms in the index case and continued for seven days. Influenza was confirmed in 163 out of 377 index cases. Oseltamivir significantly reduced the incidence of clinical influenza illness occurring in the contacts of confirmed influenza cases from 24/200 (12 %) in the placebo group to 2/205 (1 %) in the oseltamivir group (92 % reduction [95 % CI 6 – 16; $p \leq 0.0001$]). The number needed to treat (NNT) in contacts of true influenza cases was 10 (95 % CI 9 – 12) and was 16 (95 % CI 15 – 19) in the whole population (ITT) regardless of infection status in the index case.

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households that included adults, adolescents, and children aged 1 to 12 years, both as index cases and as family contacts. The primary efficacy parameter for this study was the incidence of laboratory-confirmed clinical influenza in the households. Oseltamivir prophylaxis lasted for 10 days. In the total population, there was a reduction in the incidence of laboratory-confirmed clinical influenza in households from 20 % (27/136) in the group not receiving prevention to 7 % (10/135) in the group receiving prevention (62.7 % reduction [95 % CI 26.0 – 81.2; $p = 0.0042$]). In households of influenza-infected index cases, there was a reduction in the incidence of influenza from 26 % (23/89) in the group not receiving prevention to 11 % (9/84) in the group receiving prevention (58.5 % reduction [95 % CI 15.6 – 79.6; $p = 0.0114$]). According to subgroup analysis in children at 1 to 12 years of age, the incidence of laboratory-confirmed clinical influenza among children was significantly reduced from 19 % (21/111) in the group not receiving prevention to 7 % (7/104) in the group receiving prevention (64.4 % reduction [95 % CI 15.8 – 85.0; $p = 0.0188$]). Among children who were not already shedding virus at baseline, the incidence of laboratory-confirmed clinical influenza was reduced from 21 % (15/70) in the group not receiving prevention to 4 % (2/47) in the group receiving prevention (80.1 % reduction [95 % CI 22.0 – 94.9; $p = 0.0206$]). The NNT for the total paediatric population was 9 (95 % CI 7 – 24) and 8 (95 % CI 6, upper limit not estimable) in the whole population (ITT) and in paediatric contacts of infected index cases (ITTII), respectively.

Prevention during an influenza epidemic in the community: In a pooled analysis of two other studies conducted in unvaccinated otherwise healthy adults, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 25/519 (4.8 %) in the placebo

group to 6/520 (1.2 %) in the oseltamivir group (76 % reduction [95 % CI 1.6 – 5.7; $p = 0.0006$]) during a community outbreak of influenza. The NNT in this study was 28 (95 % CI 24 – 50). A study in elderly residents of nursing homes, where 80 % of participants received vaccine in the season of the study, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 12/272 (4.4 %) in the placebo group to 1/276 (0.4 %) in the oseltamivir group (92 % reduction [95 % CI 1.5 – 6.6; $p = 0.0015$]). The NNT in this study was 25 (95 % CI 23 – 62).

Specific studies have not been conducted to assess of the reduction in the risk of complications.

5.2 Pharmacokinetic properties

Absorption

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate (pro-drug) and is extensively converted by predominantly hepatic esterases to the active metabolite (oseltamivir carboxylate). At least 75 % of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the pro-drug is less than 5 % relative to the active metabolite. Plasma concentrations of both pro-drug and active metabolite are proportional to dose and are unaffected by co-administration with food.

Distribution

The mean volume of distribution at steady state of the oseltamivir carboxylate is approximately 23 litres in humans, a volume equivalent to extracellular body fluid. Since neuraminidase activity is extracellular, oseltamivir carboxylate distributes to all sites of influenza virus spread.

The binding of the oseltamivir carboxylate to human plasma protein is negligible (approximately 3 %).

Metabolism

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. *In vitro* studies demonstrated that neither oseltamivir nor the active metabolite is a substrate for, or an inhibitor of, the major cytochrome P450 isoforms. No phase 2 conjugates of either compound have been identified *in vivo*.

Elimination

Absorbed oseltamivir is primarily (> 90 %) eliminated by conversion to oseltamivir carboxylate. It is not further metabolised and is eliminated in the urine. Peak plasma concentrations of oseltamivir carboxylate decline with a half-life of 6 to 10 hours in most subjects. The active metabolite is eliminated entirely by renal excretion. Renal clearance (18.8 l/h) exceeds glomerular filtration rate (7.5 l/h) indicating that tubular secretion occurs in addition to glomerular filtration. Less than 20 % of an oral radiolabelled dose is eliminated in faeces.

Renal impairment

Administration of 100 mg oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. For dosing, see section 4.2.

Hepatic impairment

In vitro studies have concluded that exposure to oseltamivir is not expected to be increased significantly nor is exposure to the active metabolite expected to be significantly decreased in patients with hepatic impairment (see section 4.2).

Elderly

Exposure to the active metabolite at steady state was 25 to 35 % higher in elderly (age 65 to 78 years) compared to adults less than 65 years of age given comparable doses of oseltamivir. Half-lives observed in the elderly were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage adjustments are not required for elderly patients unless there is evidence of severe renal impairment (creatinine clearance below 30 ml/min) (see section 4.2).

Children

Children 1 year of age and older: The pharmacokinetics of oseltamivir have been evaluated in single-dose pharmacokinetic studies in children aged 1 to 16 years. Multiple-dose pharmacokinetics were studied in a small number of children enrolled in a clinical efficacy study. Younger children cleared both the pro-drug and its active metabolite faster than adults, resulting in a lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in children over 12 years of age are similar to those in adults.

Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children > 1 year of age (see sections 4.1 and 4.2). There are currently no data available in infants less than 1 month of age using Tamiflu.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity and genotoxicity. Results of the conventional rodent carcinogenicity studies showed a trend towards a dose-dependent increase in the incidence of some tumours that are typical for the rodent strains used. Considering the margins of exposure in relation to the expected exposure in the human use, these findings do not change the benefit-risk of Tamiflu in its adopted therapeutic indications.

Teratology studies have been conducted in rats and rabbits at doses of up to 1500 mg/kg/day and 500 mg/kg/day, respectively. No effects on foetal development were observed. A rat fertility study up to a dose of 1500 mg/kg/day demonstrated no adverse effects on either sex. In pre- and post-natal rat studies, prolonged parturition was noted at 1500 mg/kg/day: the safety margin between human exposure and the highest no-effect dose (500 mg/kg/day) in rats is 480-fold for oseltamivir and 44-fold for the active metabolite, respectively. Foetal exposure in the rats and rabbits was approximately 15 to 20 % of that of the mother.

In lactating rats, oseltamivir and the active metabolite are excreted in the milk. Limited data indicate that oseltamivir and the active metabolite are excreted in human milk. Extrapolation of the animal data provides estimates of 0.01 mg/day and 0.3 mg/day for the respective compounds.

A potential for skin sensitisation to oseltamivir was observed in a "maximisation" test in guinea pigs. Approximately 50 % of the animals treated with the unformulated active ingredient showed erythema after challenging the induced animals. Reversible irritancy of rabbits' eyes was detected.

Whereas very high oral single doses of oseltamivir phosphate had no effect in adult rats, such doses resulted in toxicity in juvenile 7-day-old rat pups, including death. These effects were seen at doses of 657 mg/kg and higher. At 500 mg/kg, no adverse effects were seen, including upon chronic treatment (500 mg/kg/day administered from 7 to 21 days post partum).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Powder for oral suspension:

Sorbitol (E420),

Sodium dihydrogen citrate (E331[a])

Xanthan gum (E415)

Sodium benzoate (E211)

Saccharin sodium (E954)

Titanium dioxide (E171)

Tutti frutti flavour (including maltodextrins [maize], propylene glycol, arabic gum E414 and natural identical flavouring substances [mainly consisting of banana, pineapple and peach flavour]).

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

2 years for powder for oral suspension.

6.4 Special precautions for storage

Powder: Do not store above 30°C.

After reconstitution, the suspension can be stored either at room temperature (not above 25°C) for 10 days or in a refrigerator (2°C - 8°C) for 17 days.

6.5 Nature and contents of container

Carton containing a 100 ml amber glass bottle (with child-resistant plastic screw cap) with 30 g of powder for oral suspension, a plastic adapter, a plastic oral dispenser and a plastic measuring cup. After reconstitution with 52 ml of water, the usable volume of oral suspension allows for the retrieval of a total of 10 doses of 75 mg oseltamivir.

6.6 Special precautions for disposal and other handling

It is recommended that Tamiflu oral suspension should be reconstituted by the pharmacist prior to its dispensing to the patient.

Preparation of Oral Suspension

1. Tap the closed bottle gently several times to loosen the powder.
2. Measure 52 ml of water by filling the measuring cup to the indicated level (measuring cup included in the box).
3. Add all 52 ml of water into the bottle, recap the bottle and shake the closed bottle well for 15 seconds.
4. Remove the cap and push the bottle adapter into the neck of the bottle.
5. Close the bottle tightly with the cap (on the top of the bottle adapter). This will make sure that the bottle adapter fits in the bottle in the right position.

Tamiflu powder for suspension will appear as an opaque and white to light yellow suspension after reconstitution.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/02/222/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 20 June 2002
Date of last renewal: 20 June 2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>.

ANNEX II

- A. MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OF THE MARKETING AUTHORISATION**

A MANUFACTURING AUTHORISATION HOLDER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

Roche Pharma AG
Emil-Barell-Str. 1
D- 79639 Grenzach –Wyhlen
Germany

B CONDITIONS OF THE MARKETING AUTHORISATION

• **CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IMPOSED ON THE MARKETING AUTHORISATION HOLDER**

Medicinal product subject to medical prescription

• **CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

Not applicable.

• **OTHER CONDITIONS**

The MAH will continue to submit yearly PSURs (Data Lock Point 21 September) unless otherwise specified by the CHMP.

As per the CHMP Guideline on Risk Management Systems for medicinal products for human use, the updated RMP should be submitted at the same time as the next Periodic Safety Update Report (PSUR).

In addition, an updated RMP should be submitted

- When new information is received that may impact on the current Safety Specification, Pharmacovigilance Plan or risk minimisation activities.
- Within 60 days of an important (pharmacovigilance or risk minimisation) milestone being reached.
- At the request of the EMEA.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**Outer carton****1. NAME OF THE MEDICINAL PRODUCT**

Tamiflu 30 mg hard capsules
Oseltamivir

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each hard capsule contains oseltamivir phosphate equivalent to 30 mg of oseltamivir.

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

10 capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use
Read package leaflet before use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25 °C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/02/222/003

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

tamiflu 30 mg

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

Blisters

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 30 mg capsules
Oseltamivir

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Roche Registration Ltd.

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**Outer carton****1. NAME OF THE MEDICINAL PRODUCT**

Tamiflu 45 mg hard capsules
Oseltamivir

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each hard capsule contains oseltamivir phosphate equivalent to 45 mg of oseltamivir.

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

10 capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use
Read package leaflet before use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25 °C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/02/222/004

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

tamiflu 45 mg

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

Blisters

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 45 mg capsules
Oseltamivir

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Roche Registration Ltd.

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**Outer carton****1. NAME OF THE MEDICINAL PRODUCT**

Tamiflu 75 mg hard capsules
Oseltamivir

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each hard capsule contains oseltamivir phosphate equivalent to 75 mg of oseltamivir.

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

10 capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use
Read package leaflet before use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25 °C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/02/222/001

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

tamiflu 75 mg

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

Blisters

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 75 mg capsules
Oseltamivir

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Roche Registration Ltd.

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**Outer carton****1. NAME OF THE MEDICINAL PRODUCT**

Tamiflu 12 mg/ml powder for oral suspension
Oseltamivir

2. STATEMENT OF ACTIVE SUBSTANCE(S)

1 bottle with 30 g powder for oral suspension contains 900 mg of oseltamivir. Final volume of the bottle after reconstitution with 52 ml of water is 75 ml. Each ml of suspension contains 12 mg oseltamivir.

3. LIST OF EXCIPIENTS

Also contains sorbitol and sodium benzoate.

4. PHARMACEUTICAL FORM AND CONTENTS

1 bottle
Also contains 1 plastic bottle adapter, 1 plastic measuring cup (52 ml) and 1 plastic oral dispenser

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For oral administration after constitution
Shake bottle well before use
Read the enclosed leaflet before use and also for the method of preparation

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Powder: Do not store above 30°C
After reconstitution, the suspension can be stored either at room temperature (not above 25°C) for 10 days or in a refrigerator (2°C - 8°C) for 17 days

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/02/222/002

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

tamiflu

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**Label for bottle****1. NAME OF THE MEDICINAL PRODUCT**

Tamiflu 12 mg/ml powder for oral suspension
Oseltamivir

2. METHOD OF ADMINISTRATION

For oral administration after constitution
Shake bottle well before use
Read the enclosed leaflet before use and also for the method of preparation

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**6. LIST OF EXCIPIENTS**

Also contains sorbitol and sodium benzoate.

7. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children

8. SPECIAL STORAGE CONDITIONS

Powder: Do not store above 30°C
After reconstitution, the suspension can be stored either at room temperature (not above 25°C) for 10 days or in a refrigerator (2°C - 8°C) for 17 days

9. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER
--

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

B. PACKAGE LEAFLET

PACKAGE LEAFLET: INFORMATION FOR THE USER

Tamiflu 30 mg hard capsules oseltamivir

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Tamiflu is and what it is used for
2. Before you take Tamiflu
3. How to take Tamiflu
4. Possible side effects
5. How to store Tamiflu
6. Further information

1. WHAT TAMIFLU IS AND WHAT IT IS USED FOR

- Tamiflu is prescribed to you for treating or preventing influenza.
- Tamiflu belongs to a group of medicines named “neuraminidase inhibitors”. These medicines prevent the influenza virus from spreading inside the body and so help to ease or prevent the symptoms arising from the influenza virus infection.
- Influenza is an infection caused by the influenza virus. The signs (symptoms) of influenza include the sudden onset of fever (more than 37.8 °C), cough, runny or stuffy nose, headaches, muscle aches and often extreme fatigue. These symptoms may also be caused by infections other than influenza. True influenza infection only occurs during annual outbreaks (epidemics) at times when influenza viruses are spreading in the local community. Outside of the epidemic period, these symptoms will mainly be caused by a different type of infection or illness.

2. BEFORE YOU TAKE TAMIFLU

Do not take Tamiflu

- if you are allergic (hypersensitive) to oseltamivir or any of the other ingredients of Tamiflu.

Take special care with Tamiflu

Before you take Tamiflu, make sure your prescribing doctor knows if you

- are allergic to other medicines
- have problems with your kidneys.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Tamiflu can be taken with paracetamol, ibuprofen or acetylsalicylic acid (Aspirin). Tamiflu is not expected to alter the effect of any other medicines.

Are there any interactions with an influenza vaccination?

Tamiflu is not a substitute for influenza vaccination. Tamiflu will not change the effectiveness of the influenza vaccine. Even if a vaccination against influenza has been given to you Tamiflu may be prescribed by your doctor.

Taking Tamiflu with food and drink

Swallow Tamiflu with water. Tamiflu can be taken with or without food, although it is recommended to take Tamiflu with food to reduce the chance of feeling or being sick (nausea or vomiting).

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

You must tell your doctor if you are pregnant, if you think you are pregnant or if you are trying to get pregnant so that your doctor can decide if Tamiflu is right for you.

The effects on nursing infants are unknown. You must tell your doctor if you are breast-feeding so that your doctor can decide if Tamiflu is right for you.

Driving and using machines

Tamiflu has no effect on your ability to drive or use machines.

3. HOW TO TAKE TAMIFLU

Always take Tamiflu exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Take Tamiflu as soon as you get the prescription as this will help to slow the spread of the influenza virus in the body.

Swallow Tamiflu capsules whole with water. Do not break or chew Tamiflu capsules.

The usual dose is as follows:

<i>Treatment</i>

➤ Adolescents (13 to 17 years of age) and adults:

30 mg and 45 mg capsules can be used by adults and adolescents as an alternative to 75 mg capsules. For treatment of influenza take one 30 mg and one 45 mg capsule as soon as you get the prescription and then take one 30 mg and one 45 mg capsule twice a day (usually it is convenient to take one 30 mg and one 45 mg in the morning and one 30 mg and one 45 mg in the evening for five days). It is important to complete the whole 5 day course, even if you start to feel better quickly.

➤ Infants 1 year of age and older, and children 2 to 12 years of age: Tamiflu oral suspension can be used instead of capsules.

Children weighing more than 40 kg and who can swallow capsules may take a 75 mg dose of Tamiflu capsules twice daily for 5 days.

You must give the amount of capsules to your child as prescribed by your doctor.

The usual dose for treatment of influenza in children depends on the body weight of the child (see table below).

Children 1 year of age and older:

Body Weight	Recommended dose for 5 days (Treatment)
Less than or equal to 15 kg	30 mg twice daily
More than 15 kg and up to 23 kg	45 mg twice daily
More than 23 kg and up to 40 kg	60 mg twice daily
More than 40 kg	75 mg twice daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

➤ Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children older than 1 year of age (see table below for weight-based dosing recommendations). There are currently no data available in patients less than 1 month of age using Tamiflu.

Age	Recommended dose for 5 days (Treatment)
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month	2 mg/kg twice daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention

Tamiflu can also be used for prevention of influenza following exposure to an infected individual, such as family members.

➤ Adolescents (13 to 17 years of age) and adults:
30 mg and 45 mg capsules can be used by adults and adolescents as an alternative to 75 mg capsules. Tamiflu should be taken once daily for 10 days. It is best to take this dose in the mornings with breakfast.

➤ Infants 1 year of age and older, and children 2 to 12 years of age: Tamiflu oral suspension can be used instead of capsules.

The usual dose for prevention of influenza in children depends on the body weight of the child (see table below).

Children 1 year of age and older:

Body Weight	Recommended dose for 10 days (Prevention)
Less than or equal to 15 kg	30 mg once daily
More than 15 kg and up to 23 kg	45 mg once daily
More than 23 kg and up to 40 kg	60 mg once daily
More than 40 kg	75 mg once daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

Your doctor will recommend the length of time to continue taking Tamiflu, if it is prescribed to prevent influenza.

➤ Infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months of age during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children older than 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days (Prevention)
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month	2 mg/kg once daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

When Tamiflu oral suspension is not available

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Preparing Tamiflu doses for infants younger than 1 year — 30 mg capsules

Children who weigh 10 kg or less need a Tamiflu dose of 30 mg or less. To make doses of Tamiflu for these children, you will need to open a Tamiflu capsule, prepare a powder-water mixture and mix the correct volume of the mixture with food.


You will need:

1. One 30 mg Tamiflu capsule
2. Dosing chart to determine the correct dosage amount (see Tamiflu dosing chart under step 3 in the next section below)
3. A pair of scissors
4. Two small bowls per child (use separate pairs of bowls for each child)
5. One 3 ml (three millilitre) oral syringe that shows measurements of 0.1 ml (one-tenth of one millilitre)
6. One suitable oral dosing syringe that shows measurements of 0.1 ml (one-tenth of one millilitre). Ask the pharmacist for the correct syringe (1 ml, 2 ml, or 3 ml).
7. One teaspoon
8. One of these foods to hide the bitter taste of the Tamiflu powder:
 - Sugar water (If you plan to use sugar water, make this mixture now in a separate bowl or cup. Mix a teaspoon of water with three-fourths $[3/4]$ of a teaspoon of sugar.)
 - Chocolate syrup
 - Cherry syrup
 - Dessert toppings, like caramel or fudge sauce

Making the Tamiflu 10 mg/ml mixture for infants younger than 1 year — 30 mg capsules


These instructions show you how to make one dose of Tamiflu, to be used immediately.

1. Carefully open ONE 30 mg Tamiflu capsule over the first small bowl. To open a capsule, hold the capsule upright and snip off the rounded tip with scissors (see accompanying picture). Pour all of the powder into the bowl. Handle the powder carefully, because it may be irritating to the skin and eyes.



2. Follow the chart below. Use the 3 ml dosing syringe to measure 3.0 ml of water, and add it to the powder in the bowl. *Stir for about 2 minutes.*

Tamiflu capsule	Amount of water to add
ONE 30 mg capsule 	3.0 ml (three millilitres)




3. Follow the chart below. Find the child's weight on the left side of the chart, and then look at the column on the right side of chart to see the amount of the Tamiflu powder-water mixture to draw up into the dosing syringe. Draw up the correct amount of powder-water mixture from the first small bowl into the dosing syringe, and then gently expel it from the dosing syringe into the second bowl.

Tamiflu dosing chart for infants 1 to 12 months of age:

Weight (rounded to the nearest 0.5 kg)	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
4 kg	10 mg	1.00 ml
4.5 kg	11.25 mg	1.10 ml
5 kg	12.5 mg	1.30 ml
5.5 kg	13.75 mg	1.40 ml
6 kg	15 mg	1.50 ml
7 kg	21 mg	2.10 ml
8 kg	24 mg	2.40 ml
9 kg	27 mg	2.70 ml
≥ 10 kg	30 mg	3.00 ml

Tamiflu dosing chart for infants below 1 month of age:

Weight (rounded to the nearest 0.5 kg)	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
3 kg	6 mg	0.60 ml
3.5 kg	7 mg	0.70 ml
4 kg	8 mg	0.80 ml
4.5 kg	9 mg	0.90 ml



4. Add a small amount of one of the sweet foods (use no more than 1 teaspoon) into the second bowl. This will hide the bitter taste of the powder-water mixture.



5. Mix the sweet food and powder-water mixture well.



6. Give the entire contents of the second bowl (sweet food with powder-water mixture) to the child.
7. Give the child something suitable to drink.
8. Throw out any unused powder-water mixture left in the first bowl.

(Note: Not all capsule contents may dissolve. Do not worry, as the undissolved materials are inactive ingredients.)

If you take more Tamiflu, than you should

Contact your doctor or pharmacist immediately.

If you have forgotten to take Tamiflu

Do not take a double dose to make up for a forgotten capsule.

If you stop taking Tamiflu

There are no side effects when Tamiflu is discontinued prior to advice from your doctor. If Tamiflu is stopped earlier than your doctor told you, the symptoms of influenza may reoccur.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Tamiflu can cause side effects, although not everybody gets them.

- Common side effects of Tamiflu

The most common side effects of Tamiflu are nausea, vomiting, diarrhoea, stomach ache and headache. These side effects mostly occur only after the first dose of the medicine and will usually stop as treatment continues. The frequency of these effects is reduced if the medicinal product is taken with food.

- Less common side effects of Tamiflu

Adults and adolescents (children aged 13 years and older)

Other less common side effects, which may also be caused by influenza, are upper abdominal fullness, bleeding in the gastrointestinal tract, bronchitis, upper respiratory tract infections, dizziness, tiredness,

sleeping difficulties, skin reactions, mild to severe liver function disorders, visual disturbances and heart rhythm abnormalities.

Influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

During Tamiflu treatment, events like convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares) have been reported, in a very few cases resulting in accidental injury, in some instances with fatal outcome. These events were reported primarily among children and adolescents and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Children (aged 1 to 12 years)

Other less common side effects, which may also be caused by influenza, are ear inflammation, inflammation of the lungs, sinusitis, bronchitis, aggravation of pre-existing asthma, nose bleeding, ear disorders, inflammation of the skin, swelling of the lymph nodes, conjunctivitis, visual disturbances and heart rhythm abnormalities.

Infants (aged 6 to 12 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 6 to 12 months of age are similar to the side effects reported for older children (1 year and older). Please read the section above for your information.

Infants (aged 0 to 6 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 1 to 6 months of age are similar to the side effects reported for infants 6 to 12 months of age and older children (1 year and older). There are no data available for the use of Tamiflu in infants less than 1 month of age. Please read the section above for your information.

If you or your child are often sick, you should inform your doctor. You should also tell your doctor if the influenza symptoms get worse or the fever continues.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, tell your doctor or pharmacist.

5. HOW TO STORE TAMIFLU

Keep out of the reach and sight of children.

Do not use Tamiflu after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Do not store above 25 °C.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Tamiflu contains

- The active substance is oseltamivir phosphate (30 mg oseltamivir in each hard capsule).
- The other ingredients are:
capsule contents: pregelatinised starch, talc, povidone, croscarmellose sodium and sodium stearyl fumarate
capsule shell: gelatin, yellow iron oxide (E172), red iron oxide (E172) and titanium dioxide (E171)
printing ink: shellac (E904), titanium dioxide (E171) and FD and C Blue 2 (indigocarmine, E132).

What Tamiflu looks like and contents of the pack

30 mg hard capsule consists of a light yellow opaque body bearing the imprint “ROCHE” and a light yellow opaque cap bearing the imprint “30 mg”. Imprints are blue.

Tamiflu 30 mg hard capsules are available in blister packs of 10.

Marketing Authorisation Holder and Manufacturer

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

Roche Pharma AG
Emil-Barell-Str. 1,
D-79639 Grenzach-Wyhlen
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

N.V. Roche S.A.
Tél/Tel: +32 (0) 2 525 82 11

Luxembourg/Luxemburg

(Voir/siehe Belgique/Belgien)

България

Рош България ЕООД
Тел: +359 2 818 44 44

Magyarország

Roche (Magyarország) Kft.
Tel: +36 - 23 446 800

Česká republika

Roche s. r. o.
Tel: +420 - 2 20382111

Malta

(See United Kingdom)

Danmark

Roche a/s
Tlf: +45 - 36 39 99 99

Nederland

Roche Nederland B.V.
Tel: +31 (0) 348 438050

Deutschland

Roche Pharma AG
Tel: +49 (0) 7624 140

Eesti

Roche Eesti OÜ
Tel: + 372 - 6 177 380

Ελλάδα

Roche (Hellas) A.E.
Τηλ: +30 210 61 66 100

España

Roche Farma S.A.
Tel: +34 - 91 324 81 00

France

Roche
Tél: +33 (0) 1 46 40 50 00

Ireland

Roche Products (Ireland) Ltd.
Tel: +353 (0) 1 469 0700

Ísland

Roche a/s
c/o Icepharma hf
Sími: +354 540 8000

Italia

Roche S.p.A.
Tel: +39 - 039 2471

Κύπρος

Γ.Α.Σταμάτης & Σια Ατδ.
Τηλ: +357 - 22 76 62 76

Latvija

Roche Latvija SIA
Tel: +371 - 6 7039831

Lietuva

UAB "Roche Lietuva"
Tel: +370 5 2546799

Norge

Roche Norge AS
Tlf: +47 - 22 78 90 00

Österreich

Roche Austria GmbH
Tel: +43 (0) 1 27739

Polska

Roche Polska Sp.z o.o.
Tel: +48 - 22 345 18 88

Portugal

Roche Farmacêutica Química, Lda
Tel: +351 - 21 425 70 00

România

Roche România S.R.L.
Tel: +40 21 206 47 01

Slovenija

Roche farmacevtska družba d.o.o.
Tel: +386 - 1 360 26 00

Slovenská republika

Roche Slovensko, s.r.o.
Tel: +421 - 2 52638201

Suomi/Finland

Roche Oy
Puh/Tel: +358 (0) 10 554 500

Sverige

Roche AB
Tel: +46 (0) 8 726 1200

United Kingdom

Roche Products Ltd.
Tel: +44 (0) 1707 366000

This leaflet was last approved in {MM/YYYY}.

Detailed information on this medicine is available on the European Medicines Agency (EMA) web site: <http://www.ema.europa.eu/>.

PACKAGE LEAFLET: INFORMATION FOR THE USER

Tamiflu 45 mg hard capsules oseltamivir

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Tamiflu is and what it is used for
2. Before you take Tamiflu
3. How to take Tamiflu
4. Possible side effects
5. How to store Tamiflu
6. Further information

1. WHAT TAMIFLU IS AND WHAT IT IS USED FOR

- Tamiflu is prescribed to you for treating or preventing influenza.
- Tamiflu belongs to a group of medicines named “neuraminidase inhibitors”. These medicines prevent the influenza virus from spreading inside the body and so help to ease or prevent the symptoms arising from the influenza virus infection.
- Influenza is an infection caused by the influenza virus. The signs (symptoms) of influenza include the sudden onset of fever (more than 37.8 °C), cough, runny or stuffy nose, headaches, muscle aches and often extreme fatigue. These symptoms may also be caused by infections other than influenza. True influenza infection only occurs during annual outbreaks (epidemics) at times when influenza viruses are spreading in the local community. Outside of the epidemic period, these symptoms will mainly be caused by a different type of infection or illness.

2. BEFORE YOU TAKE TAMIFLU

Do not take Tamiflu

- if you are allergic (hypersensitive) to oseltamivir or any of the other ingredients of Tamiflu.

Take special care with Tamiflu

Before you take Tamiflu, make sure your prescribing doctor knows if you

- are allergic to other medicines
- have problems with your kidneys.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Tamiflu can be taken with paracetamol, ibuprofen or acetylsalicylic acid (Aspirin). Tamiflu is not expected to alter the effect of any other medicines.

Are there any interactions with an influenza vaccination?

Tamiflu is not a substitute for influenza vaccination. Tamiflu will not change the effectiveness of the influenza vaccine. Even if a vaccination against influenza has been given to you Tamiflu may be prescribed by your doctor.

Taking Tamiflu with food and drink

Swallow Tamiflu with water. Tamiflu can be taken with or without food, although it is recommended to take Tamiflu with food to reduce the chance of feeling or being sick (nausea or vomiting).

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

You must tell your doctor if you are pregnant, if you think you are pregnant or if you are trying to get pregnant so that your doctor can decide if Tamiflu is right for you.

The effects on nursing infants are unknown. You must tell your doctor if you are breast-feeding so that your doctor can decide if Tamiflu is right for you.

Driving and using machines

Tamiflu has no effect on your ability to drive or use machines.

3. HOW TO TAKE TAMIFLU

Always take Tamiflu exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Take Tamiflu as soon as you get the prescription as this will help to slow the spread of the influenza virus in the body.

Swallow Tamiflu capsules whole with water. Do not break or chew Tamiflu capsules.

The usual dose is as follows:

<i>Treatment</i>

➤ Adolescents (13 to 17 years of age) and adults:

30 mg and 45 mg capsules can be used by adults and adolescents as an alternative to 75 mg capsules. For treatment of influenza take one 30 mg and one 45 mg capsule as soon as you get the prescription and then take one 30 mg and one 45 mg capsule twice a day (usually it is convenient to take one 30 mg and one 45 mg in the morning and one 30 mg and one 45 mg in the evening for five days). It is important to complete the whole 5 day course, even if you start to feel better quickly.

➤ Infants 1 year of age and older, and children 2 to 12 years of age: Tamiflu oral suspension can be used instead of capsules.

Children weighing more than 40 kg and who can swallow capsules may take a 75 mg dose of Tamiflu capsules twice daily for 5 days.

You must give the amount of capsules to your child as prescribed by your doctor.

The usual dose for treatment of influenza in children depends on the body weight of the child (see table below).

Children 1 year of age and older:

Body Weight	Recommended dose for 5 days (Treatment)
Less than or equal to 15 kg	30 mg twice daily
More than 15 kg and up to 23 kg	45 mg twice daily
More than 23 kg and up to 40 kg	60 mg twice daily
More than 40 kg	75 mg twice daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

➤ Infants below 12 months of age: Limited pharmacokinetic and safety data are available for infants less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children older than 1 year of age (see table below for weight-based dosing recommendations). There are currently no data available in infants less than 1 month of age using Tamiflu.

Age	Recommended dose for 5 days (Treatment)
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month	2 mg/kg twice daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention

Tamiflu can also be used for prevention of influenza following exposure to an infected individual, such as family members.

➤ Adolescents (13 to 17 years of age) and adults:
30 mg and 45 mg capsules can be used by adults and adolescents as an alternative to 75 mg capsules. Tamiflu should be taken once daily for 10 days. It is best to take this dose in the mornings with breakfast.

➤ Infants 1 year of age and older, and children 2 to 12 years of age: Tamiflu oral suspension can be used instead of capsules.

The usual dose for prevention of influenza in children depends on the body weight of the child (see table below).

Children 1 year of age and older:

Body Weight	Recommended dose for 10 days (Prevention)
Less than or equal to 15 kg	30 mg once daily
More than 15 kg and up to 23 kg	45 mg once daily
More than 23 kg and up to 40 kg	60 mg once daily
More than 40 kg	75 mg once daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

Your doctor will recommend the length of time to continue taking Tamiflu, if it is prescribed to prevent influenza.

➤ Infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months of age during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children older than 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days (Prevention)
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month	2 mg/kg once daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

When Tamiflu oral suspension is not available

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Preparing Tamiflu doses for infants younger than 1 year — 45 mg capsules

Infants who weigh 10 kg or less need a Tamiflu dose of 30 mg or less. To make doses of Tamiflu for these infants, you will need to open a Tamiflu capsule, prepare a powder-water mixture and mix the correct volume of the mixture with food.


You will need:

1. One 45 mg Tamiflu capsule
2. Dosing chart to determine the correct dosage amount (see Tamiflu dosing chart under step 3 in the next section below)
3. A pair of scissors
4. Two small bowls per child (use separate pairs of bowls for each child)
5. One 5 ml (five millilitre) oral syringe that shows measurements of 0.2 ml (two-tenths of one millilitre)
6. One suitable oral dosing syringe that shows measurements of 0.1 ml (one-tenth of one millilitre). Ask the pharmacist for the correct syringe (1 ml, 2 ml, or 3 ml).
7. One teaspoon
8. One of these foods to hide the bitter taste of the Tamiflu powder:
 - Sugar water (If you plan to use sugar water, make this mixture now in a separate bowl or cup. Mix a teaspoon of water with three-fourths $[3/4]$ of a teaspoon of sugar.)
 - Chocolate syrup
 - Cherry syrup
 - Dessert toppings, like caramel or fudge sauce

Making the Tamiflu 10 mg/ml mixture for infants younger than 1 year — 45 mg capsules


These instructions show you how to make one dose of Tamiflu, to be used immediately.

1. Carefully open ONE 45 mg Tamiflu capsule over the first small bowl. To open a capsule, hold the capsule upright and snip off the rounded tip with scissors (see accompanying picture). Pour all of the powder into the bowl. Handle the powder carefully, because it may be irritating to the skin and eyes



2. Follow the chart below. Use the 5 ml dosing syringe to measure 4.5 ml of water, and add it to the powder in the bowl. *Stir for about 2 minutes.*

Tamiflu capsule	Amount of water to add
ONE 45 mg capsule 	4.5 ml (four and a half millilitres)




3. Follow the chart below. Find the child's weight on the left side of the chart, and then look at the column on the right side of chart to see the amount of the Tamiflu powder-water mixture to draw up into the dosing syringe. Draw up the correct amount of powder-water mixture from the first small bowl into the dosing syringe, and then gently expel it from the dosing syringe into the second bowl.

Tamiflu dosing chart for infants 1 to 12 months of age:

Weight (rounded to the nearest 0.5 kg)	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
4 kg	10 mg	1.00 ml
4.5 kg	11.25 mg	1.10 ml
5 kg	12.5 mg	1.30 ml
5.5 kg	13.75 mg	1.40 ml
6 kg	15 mg	1.50 ml
7 kg	21 mg	2.10 ml
8 kg	24 mg	2.40 ml
9 kg	27 mg	2.70 ml
≥ 10 kg	30 mg	3.00 ml

Tamiflu dosing chart for infants below 1 month of age:

Weight	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
3 kg	6 mg	0.60 ml
3.5 kg	7 mg	0.70 ml
4 kg	8 mg	0.80 ml
4.5 kg	9 mg	0.90 ml



4. Add a small amount of one of the sweet foods (use no more than 1 teaspoon) into the second bowl. This will hide the bitter taste of the powder-water mixture.



5. Mix the sweet food and powder-water mixture well.



6. Give the entire contents of the second bowl (sweet food with powder-water mixture) to the child.
7. Give the child something suitable to drink.
8. Throw out any unused powder-water mixture left in the first bowl.

(Note: Not all capsule contents may dissolve. Do not worry, as the undissolved materials are inactive ingredients.)

If you take more Tamiflu, than you should

Contact your doctor or pharmacist immediately.

If you have forgotten to take Tamiflu

Do not take a double dose to make up for a forgotten capsule.

If you stop taking Tamiflu

There are no side effects when Tamiflu is discontinued prior to advice from your doctor. If Tamiflu is stopped earlier than your doctor told you, the symptoms of influenza may reoccur.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Tamiflu can cause side effects, although not everybody gets them.

- Common side effects of Tamiflu

The most common side effects of Tamiflu are nausea, vomiting, diarrhoea, stomach ache and headache. These side effects mostly occur only after the first dose of the medicine and will usually stop as treatment continues. The frequency of these effects is reduced if the medicinal product is taken with food.

- Less common side effects of Tamiflu

Adults and adolescents (children aged 13 years and older)

Other less common side effects, which may also be caused by influenza, are upper abdominal fullness, bleeding in the gastrointestinal tract, bronchitis, upper respiratory tract infections, dizziness, tiredness, sleeping difficulties, skin reactions, mild to severe liver function disorders, visual disturbances and heart rhythm abnormalities.

Influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

During Tamiflu treatment, events like convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety,

nightmares) have been reported, in a very few cases resulting in accidental injury, in some instances with fatal outcome. These events were reported primarily among children and adolescents and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Children (aged 1 to 12 years)

Other less common side effects, which may also be caused by influenza, are ear inflammation, inflammation of the lungs, sinusitis, bronchitis, aggravation of pre-existing asthma, nose bleeding, ear disorders, inflammation of the skin, swelling of the lymph nodes, conjunctivitis, visual disturbances and heart rhythm abnormalities.

Infants (aged 6 to 12 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 6 to 12 months of age are similar to the side effects reported for older children (1 year and older). Please read the section above for your information.

Infants (aged 0 to 6 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 1 to 6 months of age are similar to the side effects reported for infants 6 to 12 months of age and older children (1 year and older). There are no data available for the use of Tamiflu in infants less than 1 month of age. Please read the section above for your information.

If you or your child are often sick, you should inform your doctor. You should also tell your doctor if the influenza symptoms get worse or the fever continues.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, tell your doctor or pharmacist.

5. HOW TO STORE TAMIFLU

Keep out of the reach and sight of children.

Do not use Tamiflu after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Do not store above 25 °C.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Tamiflu contains

- The active substance is oseltamivir phosphate (45 mg oseltamivir in each hard capsule).
- The other ingredients are:
capsule contents: pregelatinised starch, talc, povidone, croscarmellose sodium and sodium stearyl fumarate
capsule shell: gelatin, black iron oxide (E172) and titanium dioxide (E171)
printing ink: shellac (E904), titanium dioxide (E171) and FD and C Blue 2 (indigocarmine, E132).

What Tamiflu looks like and contents of the pack

45 mg hard capsule consists of a grey opaque body bearing the imprint “ROCHE” and a grey opaque cap bearing the imprint “45 mg”. Imprints are blue.

Tamiflu 45 mg hard capsules are available in blister packs of 10.

Marketing Authorisation Holder and Manufacturer

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

Roche Pharma AG
Emil-Barell-Str. 1,
D-79639 Grenzach-Wyhlen
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

N.V. Roche S.A.
Tél/Tel: +32 (0) 2 525 82 11

Luxembourg/Luxemburg

(Voir/siehe Belgique/Belgien)

България

Рош България ЕООД
Тел: +359 2 818 44 44

Magyarország

Roche (Magyarország) Kft.
Tel: +36 - 23 446 800

Česká republika

Roche s. r. o.
Tel: +420 - 2 20382111

Malta

(See United Kingdom)

Danmark

Roche a/s
Tlf: +45 - 36 39 99 99

Nederland

Roche Nederland B.V.
Tel: +31 (0) 348 438050

Deutschland

Roche Pharma AG
Tel: +49 (0) 7624 140

Norge

Roche Norge AS
Tlf: +47 - 22 78 90 00

Eesti

Roche Eesti OÜ
Tel: + 372 - 6 177 380

Österreich

Roche Austria GmbH
Tel: +43 (0) 1 27739

Ελλάδα

Roche (Hellas) A.E.
Τηλ: +30 210 61 66 100

Polska

Roche Polska Sp.z o.o.
Tel: +48 - 22 345 18 88

España

Roche Farma S.A.
Tel: +34 - 91 324 81 00

Portugal

Roche Farmacêutica Química, Lda
Tel: +351 - 21 425 70 00

France

Roche

Tél: +33 (0) 1 46 40 50 00

Ireland

Roche Products (Ireland) Ltd.

Tel: +353 (0) 1 469 0700

Ísland

Roche a/s

c/o Icepharma hf

Sími: +354 540 8000

Italia

Roche S.p.A.

Tel: +39 - 039 2471

Κύπρος

Γ.Α.Σταμάτης & Σια Λτδ.

Τηλ: +357 - 22 76 62 76

Latvija

Roche Latvija SIA

Tel: +371 - 6 7039831

Lietuva

UAB “Roche Lietuva”

Tel: +370 5 2546799

România

Roche România S.R.L.

Tel: +40 21 206 47 01

Slovenija

Roche farmacevtska družba d.o.o.

Tel: +386 - 1 360 26 00

Slovenská republika

Roche Slovensko, s.r.o.

Tel: +421 - 2 52638201

Suomi/Finland

Roche Oy

Puh/Tel: +358 (0) 10 554 500

Sverige

Roche AB

Tel: +46 (0) 8 726 1200

United Kingdom

Roche Products Ltd.

Tel: +44 (0) 1707 366000

This leaflet was last approved in {MM/YYYY}.

Detailed information on this medicine is available on the European Medicines Agency (EMA) web site: <http://www.ema.europa.eu/>.

PACKAGE LEAFLET: INFORMATION FOR THE USER

Tamiflu 75 mg hard capsules oseltamivir

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Tamiflu is and what it is used for
2. Before you take Tamiflu
3. How to take Tamiflu
4. Possible side effects
5. How to store Tamiflu
6. Further information

1. WHAT TAMIFLU IS AND WHAT IT IS USED FOR

- Tamiflu is prescribed to you for treating or preventing influenza.
- Tamiflu belongs to a group of medicines named “neuraminidase inhibitors”. These medicines prevent the influenza virus from spreading inside the body and so help to ease or prevent the symptoms arising from the influenza virus infection.
- Influenza is an infection caused by the influenza virus. The signs (symptoms) of influenza include the sudden onset of fever (more than 37.8 °C), cough, runny or stuffy nose, headaches, muscle aches and often extreme fatigue. These symptoms may also be caused by infections other than influenza. True influenza infection only occurs during annual outbreaks (epidemics) at times when influenza viruses are spreading in the local community. Outside of the epidemic period, these symptoms will mainly be caused by a different type of infection or illness.

2. BEFORE YOU TAKE TAMIFLU

Do not take Tamiflu

- if you are allergic (hypersensitive) to oseltamivir or any of the other ingredients of Tamiflu.

Take special care with Tamiflu

Before you take Tamiflu, make sure your prescribing doctor knows if you

- are allergic to other medicines
- have problems with your kidneys.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Tamiflu can be taken with paracetamol, ibuprofen or acetylsalicylic acid (Aspirin). Tamiflu is not expected to alter the effect of any other medicines.

Are there any interactions with an influenza vaccination?

Tamiflu is not a substitute for influenza vaccination. Tamiflu will not change the effectiveness of the influenza vaccine. Even if a vaccination against influenza has been given to you Tamiflu may be prescribed by your doctor.

Taking Tamiflu with food and drink

Swallow Tamiflu with water. Tamiflu can be taken with or without food, although it is recommended to take Tamiflu with food to reduce the chance of feeling or being sick (nausea or vomiting).

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

You must tell your doctor if you are pregnant, if you think you are pregnant or if you are trying to get pregnant so that your doctor can decide if Tamiflu is right for you.

The effects on nursing infants are unknown. You must tell your doctor if you are breast-feeding so that your doctor can decide if Tamiflu is right for you.

Driving and using machines

Tamiflu has no effect on your ability to drive or use machines.

3. HOW TO TAKE TAMIFLU

Always take Tamiflu exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Take Tamiflu as soon as you get the prescription as this will help to slow the spread of the influenza virus in the body.

Swallow Tamiflu capsules whole with water. Do not break or chew Tamiflu capsules.

The usual dose is as follows:

<i>Treatment</i>

➤ Adolescents (13 to 17 years of age) and adults: For treatment of influenza take one capsule as soon as you get the prescription and then take one capsule twice a day (usually it is convenient to take one in the morning and one in the evening for five days). It is important to complete the whole 5 day course, even if you start to feel better quickly.

30 mg and 45 mg capsules can be used by adults and adolescents as an alternative to 75 mg capsules.

➤ Infants 1 year of age and older, and children 2 to 12 years of age: Tamiflu oral suspension or 30 mg and 45 mg capsules can be used instead.

Children weighing more than 40 kg and who can swallow capsules may take Tamiflu 75 mg capsules twice daily for 5 days.

➤ Infants below 12 months of age: Limited pharmacokinetic and safety data are available for children less than 2 years of age. Pharmacokinetic modeling was undertaken using these data in addition to data from studies in adults and children older than 1 year of age. The results demonstrate that doses of 3 mg/kg twice daily for infants aged 3 to 12 months and 2.5 mg/kg twice daily for infants aged between 1 and 3 months provide exposures similar to those shown to be clinically efficacious in adults and children older than 1 year of age (see table below for weight-based dosing recommendations). There are currently no data available in infants less than 1 month of age using Tamiflu.

Age	Recommended dose for 5 days (Treatment)
> 3 months to 12 months	3 mg/kg twice daily
> 1 month to 3 months	2.5 mg/kg twice daily
0 to 1 month	2 mg/kg twice daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of treatment versus any potential risk to the infant.

Prevention

Tamiflu can also be used for prevention of influenza following exposure to an infected individual, such as family members.

➤ Adolescents (13 to 17 years of age) and adults: When used to prevent influenza following exposure to an infected individual, such as family members, Tamiflu should be taken once daily for 10 days. It is best to take this dose in the mornings with breakfast.

➤ Infants 1 year of age and older and children 2 to 12 years of age: Tamiflu oral suspension or 30 mg and 45 mg capsules can be used instead.

Children weighing more than 40 kg and can swallow capsules may take Tamiflu 75 mg capsules once daily for 10 days.

Your doctor will recommend the length of time to continue taking Tamiflu, if it is prescribed to prevent influenza.

➤ Infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months of age during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children older than 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days (Prevention)
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month	2 mg/kg once daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

When Tamiflu oral suspension is not available

During situations when commercially manufactured Tamiflu oral suspension is not readily available, adults, adolescents or children who are unable to swallow capsules may receive appropriate doses of Tamiflu by opening capsules and pouring the contents of capsules into a suitable, small amount (1 teaspoon maximum) of sweetened food product such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste. The mixture should be stirred and the entire contents given to the patient. The mixture must be swallowed immediately after its preparation.

Preparing Tamiflu doses for infants younger than 1 year — 75 mg capsules

Infants who weigh 10 kg or less need a Tamiflu dose of 30 mg or less. To make doses of Tamiflu for these infants, you will need to open a Tamiflu capsule and mix the powder inside the capsule with food.


You will need:

1. One 75 mg Tamiflu capsule
2. Dosing chart to determine the correct dosage amount (see Tamiflu dosing chart under step 3 in the next section below)
3. A pair of scissors
4. Two small bowls per child (use separate pairs of bowls for each child)
5. One 5 ml (five millilitre) oral syringe that shows measurements of 0.2 ml (two-tenths of one millilitre)
6. One suitable oral dosing syringe that shows measurements of 0.1 ml (one-tenth of one millilitre). Ask the pharmacist for the correct syringe (1 ml, 2 ml, or 3 ml).
7. One teaspoon
8. One of these foods to hide the bitter taste of the Tamiflu powder:
 - Sugar water (If you plan to use sugar water, make this mixture now in a separate bowl or cup. Mix a teaspoon of water with three-fourths $[3/4]$ of a teaspoon of sugar.)
 - Chocolate syrup
 - Cherry syrup
 - Dessert toppings, like caramel or fudge sauce


Making the Tamiflu 10 mg/ml mixture for infants younger than 1 year — 75 mg capsules

These instructions show you how to make one dose of Tamiflu, to be used immediately.

1. Carefully open ONE 75 mg Tamiflu capsule over the first small bowl. To open a capsule, hold the capsule upright and snip off the rounded tip with scissors (see accompanying picture). Pour all of the powder into the bowl. Handle the powder carefully, because it may be irritating to the skin and eyes.



2. Follow the chart below. Use the 5 ml dosing syringe to measure first 5.0 ml of water, and add it to the powder in the bowl. Then measure 2.5 ml of water, and add it to the powder in the bowl. *Stir for about 2 minutes.*

Tamiflu capsule		Amount of water to add
ONE 75 mg capsule		7.5 ml (seven and a half millilitres)




3. Follow the chart below. Find the child's weight on the left side of the chart, and then look at the column on the right side of chart to see the amount of the Tamiflu powder-water mixture to draw up into the dosing syringe. Draw up the correct amount of powder-water mixture from the first small bowl into the dosing syringe, and then gently expel it from the dosing syringe into the second bowl.

Tamiflu dosing chart for infants 1 to 12 months of age:

Weight (rounded to the nearest 0.5 kg)	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
4 kg	10 mg	1.00 ml
4.5 kg	11.25 mg	1.10 ml
5 kg	12.5 mg	1.30 ml
5.5 kg	13.75 mg	1.40 ml
6 kg	15 mg	1.50 ml
7 kg	21 mg	2.10 ml
8 kg	24 mg	2.40 ml
9 kg	27 mg	2.70 ml
≥ 10 kg	30 mg	3.00 ml

Tamiflu dosing chart for infants below 1 month of age:

Weight (rounded to the nearest 0.5 kg)	Required dose of Tamiflu	Amount of Tamiflu powder-water mixture to be drawn up into the dosing syringe
3 kg	6 mg	0.60 ml
3.5 kg	7 mg	0.70 ml
4 kg	8 mg	0.80 ml
4.5 kg	9 mg	0.90 ml



4. Add a small amount of one of the sweet foods (use no more than 1 teaspoon) into the second bowl. This will hide the bitter taste of the powder-water mixture.



5. Mix the sweet food and powder-water mixture well.



6. Give the entire contents of the second bowl (sweet food with powder-water mixture) to the child.
7. Give the child something suitable to drink.
8. Throw out any unused powder-water mixture left in the first bowl.

(Note: Not all capsule contents may dissolve. Do not worry, as the undissolved materials are inactive ingredients.)

Preparing Tamiflu doses for children 1 year of age or older — if you have only 75 mg capsules available

Children who weigh less than 40 kg need a Tamiflu dose of less than 75 mg. To make doses of Tamiflu for these children, you will need to open a 75 mg capsule, prepare a powder-water mixture and mix the correct volume of the mixture with food.

You will need

1. One 75 mg Tamiflu capsule
2. Dosing chart to determine the correct dosage amount (see charts under step 3 in the next section below)
3. A pair of scissors
4. Two small bowls per child (use separate pairs of bowls for each child)
5. One 5 ml (five millilitre) oral syringe that shows measurements of 0.2 ml (two-tenths of one millilitre)
6. One suitable oral dosing syringe that shows measurements of 0.1 ml (one-tenth of one millilitre). Ask the pharmacist for the correct syringe (1 ml, 2 ml, or 3 ml).
7. One teaspoon
8. One of these foods to hide the bitter taste of the Tamiflu powder:
 - Sugar water (If you plan to use sugar water, make this mixture now in a separate bowl or cup. Mix a teaspoon of water with three-fourths $[3/4]$ of a teaspoon of sugar.)
 - Chocolate syrup
 - Cherry syrup
 - Dessert toppings, like caramel or fudge sauce

Making the Tamiflu mixture for children 1 year of age or older — if you have only 75 mg capsules available

These instructions show you how to make one dose of Tamiflu, to be used immediately.

1. Carefully open ONE 75 mg Tamiflu capsule over the first small bowl. To open a capsule, hold the capsule upright and snip off the rounded tip with scissors (see accompanying picture). Pour all of the powder into the bowl. Handle the powder carefully, because it may be irritating to the skin and eyes.



2. Use the 5 ml syringe to add 5 ml of water to the powder in the bowl. *Stir for about 2 minutes.*



(Note: Not all capsule contents may dissolve. Do not worry, as the undissolved materials are inactive ingredients.)

3. The next step will be to draw up the correct amount of powder-water mixture from the first small bowl, and to gently expel it into the second small bowl.


You must find what the correct amount of powder-water mixture to draw up into the dosing syringe is. If you know the child's weight, you can do this by finding the child's weight on the left side of the first table below. Then, look at the middle column of the table to see the correct amount of the powder-water mixture to draw up into the dosing syringe.

If you do not know the child's weight, find the child's age on the left side of the second table below. Then, look at the middle column of the table to see the correct amount of the powder-water mixture to draw up into the dosing syringe.

Weight	Amount of Tamiflu Mixture	Dose in mg
Up to 15 kg	2 ml = 1 dose	30 mg
15 kg up to 23 kg	3 ml = 1 dose	45 mg
23 kg up to 40 kg	4 ml = 1 dose	60 mg

Age	Amount of Tamiflu Mixture	Dose in mg
1 to 2 years	2 ml = 1 dose	30 mg
3 to 5 years	3 ml = 1 dose	45 mg
6 to 9 years	4 ml = 1 dose	60 mg

Once you have drawn up the correct amount of powder-water mixture into the dosing syringe, gently expel it from the dosing syringe into the second bowl.



4. Add a small amount of one of the sweet foods into the second bowl to hide the bitter taste of the powder-water mixture.



5. Mix the sweet food and powder-water mixture well.



6. Give the entire contents of the second bowl (sweet food with powder-water mixture) to the child.
7. Throw out any unused powder-water mixture left in the first bowl.

Repeat this procedure every time you need to take the medicine.

If you take more Tamiflu, than you should

Contact your doctor or pharmacist immediately.

If you have forgotten to take Tamiflu

Do not take a double dose to make up for a forgotten capsule.

If you stop taking Tamiflu

There are no side effects when Tamiflu is discontinued prior to advice from your doctor. If Tamiflu is stopped earlier than your doctor told you, the symptoms of influenza may reoccur.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Tamiflu can cause side effects, although not everybody gets them.

- Common side effects of Tamiflu

The most common side effects of Tamiflu are nausea, vomiting, diarrhoea, stomach ache and headache. These side effects mostly occur only after the first dose of the medicine and will usually stop as treatment continues. The frequency of these effects is reduced if the medicinal product is taken with food.

- Less common side effects of Tamiflu

Adults and adolescents (children aged 13 years and older)

Other less common side effects, which may also be caused by influenza, are upper abdominal fullness, bleeding in the gastrointestinal tract, bronchitis, upper respiratory tract infections, dizziness, tiredness, sleeping difficulties, skin reactions, mild to severe liver function disorders, visual disturbances and heart rhythm abnormalities.

Influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

During Tamiflu treatment, events like convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares) have been reported, in a very few cases resulting in accidental injury, in some instances with fatal outcome. These events were reported primarily among children and adolescents and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Children (aged 1 to 12 years)

Other less common side effects, which may also be caused by influenza, are ear inflammation, inflammation of the lungs, sinusitis, bronchitis, aggravation of pre-existing asthma, nose bleeding, ear disorders, inflammation of the skin, swelling of the lymph nodes, conjunctivitis, visual disturbances and heart rhythm abnormalities.

Infants (aged 6 to 12 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 6 to 12 months of age are similar to the side effects reported for older children (1 year and older). Please read the section above for your information.

Infants (aged 0 to 6 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 1 to 6 months of age are similar to the side effects reported for infants 6 to 12 months of age and older children (1 year and older). There are no data available for the use of Tamiflu in infants less than 1 month of age. Please read the section above for your information.

If you or your child are often sick, you should inform your doctor. You should also tell your doctor if the influenza symptoms get worse or the fever continues.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, tell your doctor or pharmacist.

5. HOW TO STORE TAMIFLU

Keep out of the reach and sight of children.

Do not use Tamiflu after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Do not store above 25 °C.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Tamiflu contains

- The active substance is oseltamivir phosphate (75 mg oseltamivir in each hard capsule).
- The other ingredients are:
capsule contents: pregelatinised starch, talc, povidone, croscarmellose sodium and sodium stearyl fumarate
capsule shell: gelatin, yellow iron oxide (E172), red iron oxide (E172), black iron oxide (E172) and titanium dioxide (E171)
printing ink: shellac (E904), titanium dioxide (E171) and indigocarmine (E132).

What Tamiflu looks like and contents of the pack

The hard capsule consists of a grey opaque body bearing the imprint “ROCHE” and a light yellow opaque cap bearing the imprint “75 mg”. Imprints are blue.

Tamiflu 75 mg hard capsules are available in blister packs of 10.

Marketing Authorisation Holder and Manufacturer

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

Roche Pharma AG
Emil-Barell-Str. 1,
D-79639 Grenzach-Wyhlen
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

N.V. Roche S.A.

Tél/Tel: +32 (0) 2 525 82 11

България

Рош България ЕООД

Тел: +359 2 818 44 44

Česká republika

Roche s. r. o.

Tel: +420 - 2 20382111

Danmark

Roche a/s

Tlf: +45 - 36 39 99 99

Deutschland

Roche Pharma AG

Tel: +49 (0) 7624 140

Eesti

Roche Eesti OÜ

Tel: + 372 - 6 177 380

Ελλάδα

Roche (Hellas) A.E.

Τηλ: +30 210 61 66 100

España

Roche Farma S.A.

Tel: +34 - 91 324 81 00

France

Roche

Tél: +33 (0) 1 46 40 50 00

Ireland

Roche Products (Ireland) Ltd.

Tel: +353 (0) 1 469 0700

Ísland

Roche a/s

c/o Icepharma hf

Sími: +354 540 8000

Italia

Roche S.p.A.

Tel: +39 - 039 2471

Κύπρος

Γ.Α.Σταμάτης & Σια Ατδ.

Τηλ: +357 - 22 76 62 76

Latvija

Roche Latvija SIA

Tel: +371 - 6 7039831

Luxembourg/Luxemburg

(Voir/siehe Belgique/Belgien)

Magyarország

Roche (Magyarország) Kft.

Tel: +36 - 23 446 800

Malta

(See United Kingdom)

Nederland

Roche Nederland B.V.

Tel: +31 (0) 348 438050

Norge

Roche Norge AS

Tlf: +47 - 22 78 90 00

Österreich

Roche Austria GmbH

Tel: +43 (0) 1 27739

Polska

Roche Polska Sp.z o.o.

Tel: +48 - 22 345 18 88

Portugal

Roche Farmacêutica Química, Lda

Tel: +351 - 21 425 70 00

România

Roche România S.R.L.

Tel: +40 21 206 47 01

Slovenija

Roche farmacevtska družba d.o.o.

Tel: +386 - 1 360 26 00

Slovenská republika

Roche Slovensko, s.r.o.

Tel: +421 - 2 52638201

Suomi/Finland

Roche Oy

Puh/Tel: +358 (0) 10 554 500

Sverige

Roche AB

Tel: +46 (0) 8 726 1200

United Kingdom

Roche Products Ltd.

Tel: +44 (0) 1707 366000

Lietuva
UAB “Roche Lietuva”
Tel: +370 5 2546799

This leaflet was last approved in {MM/YYYY}.

Detailed information on this medicine is available on the European Medicines Agency (EMA) web site: <http://www.emea.europa.eu/>.

PACKAGE LEAFLET: INFORMATION FOR THE USER

Tamiflu 12 mg/ml powder for oral suspension oseltamivir

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Tamiflu is and what it is used for
2. Before you take Tamiflu
3. How to take Tamiflu
4. Possible side effects
5. How to store Tamiflu
6. Further information

1. WHAT TAMIFLU IS AND WHAT IT IS USED FOR

- Tamiflu is prescribed to you for treating or preventing influenza.
- Tamiflu belongs to a group of medicines named “neuraminidase inhibitors”. These medicines prevent the influenza virus from spreading inside the body and so help to ease or prevent the symptoms arising from the influenza virus infection.
- Influenza is an infection caused by the influenza virus. The signs (symptoms) of influenza include the sudden onset of fever (more than 37.8 °C), cough, runny or stuffy nose, headaches, muscle aches and often extreme fatigue. These symptoms may also be caused by infections other than influenza. True influenza infection only occurs during annual outbreaks (epidemics) at times when influenza viruses are spreading in the local community. Outside of the epidemic period, these symptoms will mainly be caused by a different type of infection or illness.

2. BEFORE YOU TAKE TAMIFLU

Do not take Tamiflu

- if you are allergic (hypersensitive) to oseltamivir or any of the other ingredients of Tamiflu.

Take special care with Tamiflu

Before you take Tamiflu, make sure your prescribing doctor knows if you

- are allergic to other medicines
- have problems with your kidneys

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Tamiflu can be taken with paracetamol, ibuprofen or acetylsalicylic acid (Aspirin). Tamiflu is not expected to alter the effect of any other medicines.

Are there any interactions with an influenza vaccination?

Tamiflu is not a substitute for influenza vaccination. Tamiflu will not change the effectiveness of the influenza vaccine. Even if a vaccination against influenza has been given to you Tamiflu may be prescribed by your doctor.

Taking Tamiflu with food and drink

Tamiflu oral suspension can be taken with or without food, although it is recommended to take Tamiflu with food to reduce the chance of feeling or being sick (nausea or vomiting). It can also be taken with a drink of water.

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

You must tell your doctor if you are pregnant, if you think you are pregnant or if you are trying to get pregnant so that your doctor can decide if Tamiflu is right for you.

The effects on nursing infants are unknown. You must tell your doctor if you are breast-feeding so that your doctor can decide if Tamiflu is right for you.

Driving and using machines

Tamiflu has no effect on your ability to drive or use machines.

Important information about some of the ingredients of Tamiflu

Before you take Tamiflu, make sure your prescribing doctor knows if you have hereditary fructose intolerance.

This medicine contains sorbitol which is a form of fructose.

3. HOW TO TAKE TAMIFLU

Always take Tamiflu exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Take Tamiflu as soon as you get the prescription as this will help to slow the spread of the influenza virus in the body.

The usual dose is as follows:

How much Tamiflu should be given to infants 1 year of age and older and children 2 to 12 years of age)**Treatment**

You must give the amount of oral suspension to your child as prescribed by your doctor.

30 mg and 45 mg capsules can be used as an alternative to suspension.

The usual dose for treatment of influenza in children depends on the body weight of the child (see table below):

Body Weight	Recommended dose for 5 days (Treatment)
Less than or equal to 15 kg	30 mg twice daily
More than 15 kg and up to 23 kg	45 mg twice daily
More than 23 kg and up to 40 kg	60 mg twice daily
More than 40 kg	75 mg twice daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

Prevention

Tamiflu can also be used to prevent influenza following exposure to an infected individual, such as family members.

30 mg and 45 mg capsules can be used as an alternative to suspension.

Tamiflu should be taken once daily for 10 days. It is best to take this dose in the mornings with breakfast.

The usual dose for prevention of influenza in children depends on the body weight of the child (see table below):

Body Weight	Recommended dose for 10 days (Prevention)
Less than or equal to 15 kg	30 mg once daily
More than 15 kg and up to 23 kg	45 mg once daily
More than 23 kg and up to 40 kg	60 mg once daily
More than 40 kg	75 mg once daily (75 mg is made up of a 30 mg dose plus a 45 mg dose)

Children weighing more than 40 kg and can swallow capsules may take a 75 mg dose of Tamiflu capsules once daily for 10 days.

Your doctor will recommend the length of time to continue taking Tamiflu, if it is prescribed to prevent influenza.

➤ Infants below 12 months of age: The recommended prophylaxis dose for infants less than 12 months of age during a pandemic influenza outbreak is half of the daily treatment dose. This is based upon clinical data in children older than 1 year of age and adults showing that a prophylaxis dose equivalent to half the daily treatment dose is clinically efficacious for the prevention of influenza. The following weight-adjusted dosing prophylaxis regimens are recommended for infants below 1 year of age:

Age	Recommended dose for 10 days (Prevention)
> 3 months to 12 months	3 mg/kg once daily
> 1 month to 3 months	2.5 mg/kg once daily
0 to 1 month	2 mg/kg once daily

Administration of Tamiflu to infants less than one year of age should be based upon the judgment of the physician after considering the potential benefit of prophylaxis versus any potential risk to the infant.

How Much Tamiflu should be given to Adolescents (13 to 17 years of age) and Adults

The suspension can be used by adults and adolescents instead of the capsule. 30 mg and 45 mg capsules can be used as an alternative to suspension.

Treatment

Usually, your doctor will prescribe a 75 mg dose of Tamiflu twice daily for 5 days.

Prevention

Usually, your doctor will prescribe a 75 mg dose of Tamiflu once daily for 10 days. Your doctor will recommend the length of time you will need to take Tamiflu.

A single 75 mg dose of Tamiflu requires two doses of the suspension: one 30 mg dose and one 45 mg dose.

a) HOW TO PREPARE THE ORAL SUSPENSION:

*Your pharmacist may have prepared the oral suspension for you when you collected your prescription. However, if they have not done this, then you can do it easily yourself. **You only need to prepare the***

suspension once, at the beginning of your course. After that, all you need to do is shake the suspension well and draw up the appropriate recommended dose.




Figure 1

1. Tap the closed bottle gently several times to loosen the powder.
2. Measure 52 ml of water by filling the measuring cup to the indicated level (measuring cup included in the box). **You should always use 52 ml of water**, irrespective of the recommended dose you are taking.
3. Add all 52 ml of water into the bottle, recap the bottle and shake the closed bottle well for 15 seconds.
4. Remove the cap and push the bottle adapter into the neck of the bottle.
5. Place the cap tightly over the top of the bottle, which now includes the bottle adapter. This will make sure that the bottle adapter fits in the bottle in the right position.

b) HOW TO MEASURE AND GIVE A DOSE OF ORAL SUSPENSION

Always shake Tamiflu oral suspension well before use!




Figure 2



Figure 3

How to measure the correct amount of suspension:

1. Shake the closed bottle of Tamiflu oral suspension well before use.
2. Take the dispenser (see figure 2) and push the plunger completely down toward the tip of the dispenser. The dispenser provided with your medicine must always be used to measure to correct dose.
3. Remove the cap from the bottle of oral suspension.
4. Insert the tip of the dispenser into the bottle adapter.
5. Turn the entire unit (bottle and dispenser) upside down (see figure 3).
6. Slowly pull out the plunger to the graduation which marks the dose you need.
7. Turn the entire unit upright.
8. Slowly remove the dispenser from the bottle.
9. Push the suspension directly into the mouth by pushing down the plunger of the dispenser. Swallow the medicine. You may drink and eat something after taking the medicine.
10. Immediately after administration, take the dispenser apart and rinse both parts of the dispenser under running tap water.

If you take more Tamiflu, than you should

Contact your doctor or pharmacist immediately.

If you have forgotten to take Tamiflu

Do not take a double dose to make up for a forgotten dose.

If you stop taking Tamiflu

There are no side effects when Tamiflu is discontinued prior to advice from your doctor. If Tamiflu is stopped earlier than your doctor told you, the symptoms of influenza may reoccur.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Tamiflu can cause side effects, although not everybody gets them.

- Common side effects of Tamiflu

The most common side effects of Tamiflu are nausea, vomiting, diarrhoea, stomach ache and headache. These side effects mostly occur only after the first dose of the medicine and will usually stop as treatment continues. The frequency of these effects is reduced if the medicinal product is taken with food.

- Less common side effects of Tamiflu**Adults and adolescents (children aged 13 years and older)**

Other less common side effects, which may also be caused by influenza, are upper abdominal fullness, bleeding in the gastrointestinal tract, bronchitis, upper respiratory tract infections, dizziness, tiredness, sleeping difficulties, skin reactions, mild to severe liver function disorders, visual disturbances and heart rhythm abnormalities.

Influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

During Tamiflu treatment, events like convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares) have been reported, in a very few cases resulting in accidental injury, in some instances with fatal outcome. These events were reported primarily among children and adolescents and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Children (aged 1 to 12 years)

Other less common side effects, which may also be caused by influenza, are ear inflammation, inflammation of the lungs, sinusitis, bronchitis, aggravation of pre-existing asthma, nose bleeding, ear disorders, inflammation of the skin, swelling of the lymph nodes, conjunctivitis, visual disturbances and heart rhythm abnormalities.

Infants (aged 6 to 12 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 6 to 12 months of age are similar to the side effects reported for older children (1 year and older). Please read the section above for your information.

Infants (aged 0 to 6 months)

The reported side effects of Tamiflu when used for treatment of influenza in infants 1 to 6 months of age are similar to the side effects reported for infants 6 to 12 months of age and older children (1 year and older). There are no data available for the use of Tamiflu in infants less than 1 month of age. Please read the section above for your information.

If you or your child are often sick, you should inform your doctor. You should also tell your doctor if the influenza symptoms get worse or the fever continues.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE TAMIFLU

Keep out of the reach and sight of children.

Do not use Tamiflu powder for oral suspension after the expiry date which is stated on the carton and bottle after EXP. The expiry date refers to the last day of that month.

Powder: Do not store above 30°C.

After reconstitution, the suspension can be stored either at room temperature (not above 25°C) for 10 days or in a refrigerator (2°C - 8°C) for 17 days.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION**What Tamiflu contains**

- The active substance is oseltamivir phosphate (12 mg/ml oseltamivir after reconstitution).
- The other ingredients are sorbitol (E420), sodium dihydrogen citrate (E331[a]), xanthan gum (E415), sodium benzoate (E211), saccharin sodium (E954), titanium dioxide (E171) and flavour.

What Tamiflu looks like and contents of the pack

The powder is a granulate or clumped granulate with a white to light yellow colour.

Tamiflu 12 mg/ml powder for oral suspension is available in a bottle containing 30 g powder for mixing with 52 ml of water.

The box also contains 1 plastic measuring cup (52 ml), 1 plastic bottle adapter (to help get the drug into the dispenser) and 1 plastic oral dispenser (to give the correct amount of medicine via the mouth). Shown on the oral dispenser are marks for 30 mg, 45 mg and 60 mg of the medicine (see figures 1 and 2 above).

For details on how to prepare the oral suspension and how to measure and take the medicine, read section 3, "How to take Tamiflu".

Marketing Authorisation Holder and Manufacturer

Roche Registration Limited
6 Falcon Way
Shire Park
Welwyn Garden City
AL7 1TW
United Kingdom

Roche Pharma AG
Emil-Barell-Str. 1,
D-79639 Grenzach-Wyhlen
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

N.V. Roche S.A.
Tél/Tel: +32 (0) 2 525 82 11

Luxembourg/Luxemburg

(Voir/siehe Belgique/Belgien)

България

Рош България ЕООД
Тел: +359 2 818 44 44

Magyarország

Roche (Magyarország) Kft.
Tel: +36 - 23 446 800

Česká republika

Roche s. r. o.
Tel: +420 - 2 20382111

Malta

(See United Kingdom)

Danmark

Roche a/s
Tlf: +45 - 36 39 99 99

Nederland

Roche Nederland B.V.
Tel: +31 (0) 348 438050

Deutschland

Roche Pharma AG
Tel: +49 (0) 7624 140

Norge

Roche Norge AS
Tlf: +47 - 22 78 90 00

Eesti

Roche Eesti OÜ
Tel: + 372 - 6 177 380

Österreich

Roche Austria GmbH
Tel: +43 (0) 1 27739

Ελλάδα

Roche (Hellas) A.E.
Τηλ: +30 210 61 66 100

Polska

Roche Polska Sp.z o.o.
Tel: +48 - 22 345 18 88

España

Roche Farma S.A.
Tel: +34 - 91 324 81 00

Portugal

Roche Farmacêutica Química, Lda
Tel: +351 - 21 425 70 00

France

Roche
Tél: +33 (0) 1 46 40 50 00

România

Roche România S.R.L.
Tel: +40 21 206 47 01

Ireland

Roche Products (Ireland) Ltd.
Tel: +353 (0) 1 469 0700

Slovenija

Roche farmacevtska družba d.o.o.
Tel: +386 - 1 360 26 00

Ísland

Roche a/s
c/o Icepharma hf
Sími: +354 540 8000

Italia

Roche S.p.A.
Tel: +39 - 039 2471

Κύπρος

Γ.Α.Σταμάτης & Σια Λτδ.
Τηλ: +357 - 22 76 62 76

Latvija

Roche Latvija SIA
Tel: +371 - 6 7039831

Lietuva

UAB “Roche Lietuva”
Tel: +370 5 2546799

Slovenská republika

Roche Slovensko, s.r.o.
Tel: +421 - 2 52638201

Suomi/Finland

Roche Oy
Puh/Tel: +358 (0) 10 554 500

Sverige

Roche AB
Tel: +46 (0) 8 726 1200

United Kingdom

Roche Products Ltd.
Tel: +44 (0) 1707 366000

This leaflet was last approved in {MM/YYYY}.

Detailed information on this medicine is available on the European Medicines Agency (EMA) web site: <http://www.ema.europa.eu/>.